

## S7 Supporting Information. Species Narratives

### Table of Contents

Acadian Redfish.....	1
Alewife .....	4
American Eel .....	9
American Lobster .....	13
American Plaice.....	17
American Shad .....	20
Anchovies.....	24
Atlantic Cod.....	27
Atlantic Croaker .....	31
Atlantic Hagfish .....	35
Atlantic Halibut .....	38
Atlantic Herring.....	41
Atlantic Mackerel .....	45
Atlantic Menhaden .....	48
Atlantic Salmon .....	52
Atlantic Saury .....	56
Atlantic Sea Scallop .....	59
Atlantic Sturgeon .....	62
Atlantic Surfclam.....	66
Atlantic Wolfish.....	69
Barndoor Skate .....	73
Bay Scallop .....	76
Black Sea Bass .....	79
Bloodworm.....	82
Blue Crab.....	85
Blue Mussel.....	89
Blueback Herring.....	93
Bluefish .....	97
Butterfish .....	101

Cancer Crabs .....	104
Channeled Whelk .....	107
Clearnose Skate.....	110
Conger Eel .....	113
Cusk.....	116
Deep-sea Red Crab.....	119
Dusky Shark.....	122
Eastern Oyster.....	126
Green Sea Urchin .....	130
Haddock .....	133
Hickory Shad .....	136
Horseshoe Crab.....	139
Knobbed Whelk.....	142
Little Skate.....	145
Longfin Inshore Squid .....	148
Monkfish (Goosefish).....	152
Northern Kingfish.....	155
Northern Quahog.....	158
Northern Shortfin Squid.....	161
Northern Shrimp .....	164
Ocean Pout.....	168
Ocean Quahog .....	171
Offshore Hake .....	174
Pollock.....	177
Porbeagle .....	181
Rainbow Smelt .....	184
Red Drum .....	188
Red Hake .....	191
Rosette Skate .....	194
Sand Lances.....	197
Sand Tiger .....	200
Scup.....	203
Shortnose Sturgeon .....	206

Silver Hake.....	210
Smooth Dogfish.....	213
Smooth Skate .....	217
Softshell Clam .....	220
Spanish Mackerel .....	224
Spiny Dogfish.....	227
Spot .....	230
Spotted Seatrout.....	233
Striped Bass.....	236
Summer Flounder .....	240
Tautog .....	244
Thorny Skate .....	248
Tilefish.....	251
Weakfish .....	255
White Hake .....	258
Windowpane.....	261
Winter Flounder.....	264
Winter Skate .....	268
Witch Flounder .....	271
Yellowtail Flounder .....	274

Eastern Oyster – *Crassostrea virginica*

Overall Vulnerability Rank = Very High ■

Biological Sensitivity = High ■

Climate Exposure = Very High ■

Data Quality = 88% of scores  $\geq 2$

<i>Crassostrea virginica</i>	Expert Scores	Data Quality	Expert Scores Plots (Portion by Category)
Stock Status	2.6	1.4	
Other Stressors	3.4	3.0	
Population Growth Rate	2.5	2.0	
Spawning Cycle	1.8	3.0	
Complexity in Reproduction	2.3	2.6	
Early Life History Requirements	2.6	3.0	
Sensitivity to Ocean Acidification	3.9	2.2	
Prey Specialization	2.1	2.6	
Habitat Specialization	1.8	2.6	
Sensitivity to Temperature	1.4	3.0	
Adult Mobility	4.0	3.0	
Dispersal & Early Life History	2.3	3.0	
<b>Sensitivity Score</b>	<b>High</b>		
Sea Surface Temperature	4.0	3.0	
Variability in Sea Surface Temperature	1.0	3.0	
Salinity	2.0	3.0	
Variability Salinity	1.2	3.0	
Air Temperature	4.0	3.0	
Variability Air Temperature	1.0	3.0	
Precipitation	1.3	3.0	
Variability in Precipitation	1.4	3.0	
Ocean Acidification	4.0	2.0	
Variability in Ocean Acidification	1.0	2.2	
Currents	2.0	1.0	
Sea Level Rise	2.6	1.5	
<b>Exposure Score</b>	<b>Very High</b>		
<b>Overall Vulnerability Rank</b>	<b>Very High</b>		

### **Eastern Oyster (*Crassostrea virginica*)**

Overall Climate Vulnerability Rank: **Very High** (100% certainty from bootstrap analysis).

Climate Exposure: **Vey High**. Three exposure factors contributed to this score: Ocean Surface Temperature (4.0), Air Temperature (4.0) and Ocean Acidification (4.0). Eastern Oyster utilize near coastal habitats and have a calcium carbonate shell.

Biological Sensitivity: **High**. Three sensitivity attributes scored above 3.0: Sensitivity to Ocean Acidification (3.9), Adult Mobility (4.0) and Other Stressors (3.4). Eastern Oyster are sessile and have a calcium carbonate shell. Adults are reef building and found in estuarine habitats. They are subject to a number of other stressors including contaminants, invasive species, and habitat loss.

Distributional Vulnerability Rank: **Moderate** (93% certainty from bootstrap analysis).

Directional Effect in the Northeast U.S. Shelf: The effect of climate change on Eastern Oyster on the Northeast U.S. Shelf is very likely to be negative (>95% certainty in expert scores). Ocean acidification will likely negatively impact molluscs, including Eastern Oyster. Increased precipitation may also reduce available habitat in estuarine areas as upper estuaries become fresher.

Data Quality: 88% of the data quality scores were 2 or greater indicate that data quality is moderate.

Climate Effects on Abundance and Distribution: Shell growth of juvenile Eastern Oysters is lower under lower aragonite saturation states (Ries et al., 2009) and lower pH (Waldbusser et al., 2011). Larvae experienced lowered growth and delayed metamorphosis at higher CO<sub>2</sub> concentrations (Talmage and Goblet, 2009). Kimmel and Newell (2007) indicated that reduced recruitment in Chesapeake Bay was due to decreased spawning stock biomass and climate-driven changes in environmental conditions, mostly changes in precipitation and freshwater input into the Bay (Kimmel et al., 2012).

Life History Synopsis: Eastern Oyster is an estuarine keystone species found from the Gulf of St. Lawrence, Canada, to Key Biscayne, Florida, the Gulf of Mexico to the Yucatan Peninsula, and the West Indies to Venezuela (Sellers and Stanley, 1984). Eastern Oyster has also been introduced to Japan, Australia, Great Britain, Hawaii, and the west coast of North America (Sellers and Stanley, 1984). The species is dioecious, but most young individuals are males while older individuals are females, and there is evidence that individuals can change gender back and forth annually if needed (Sellers and Stanley, 1984; EOBRT, 2007). Spawning is stimulated by temperature, but the spawning temperature is stock-specific (Sellers and Stanley, 1984). In northern climates, spawning occurs in the summer, and can occur year round in southern climates if temperatures remain above 20°C and salinities above 10 (EOBRT, 2007). Males release gametes into the water column first, inducing females to release eggs (Sellers and Stanley, 1984). Female oysters are highly fecund, producing tens of millions of eggs, and can spawn multiple times per season (Sellers and Stanley, 1984). Eggs incubate for 4-6 hours, and larvae progress through a series of planktonic stages over a period of 2-3 weeks post-fertilization, but can delay the settlement process up to 2 months to search for suitable attachment sites (Sellers and Stanley, 1984; EOBRT, 2007). The initial larval stage, the trochophore stage, lasts 1-2 days and is non-feeding (EOBRT, 2007). The veliger stages are marked by the secretion of a shell, development of a ciliated velum for swimming and feeding on planktonic organisms (Sellers and Stanley, 1984; EOBRT, 2007). Larvae do not tolerate high temperatures and have a narrower salinity tolerance range than adults (Sellers and Stanley, 1984; EOBRT, 2007). Larvae can be quite abundant in the plankton and are food for a variety of

filter feeders (Sellers and Stanley, 1984). By the end of the veliger stage, the larvae have a foot with a byssal gland and react to changes in salinity by changing depth distribution, aiding retention within the estuary (Sellers and Stanley, 1984). At the end of the larval stage, and likely triggered by rising temperatures, young oysters drop to the bottom and begin crawling in a circular search area, seeking substrate (Sellers and Stanley, 1984). Shells seem to be the preferred substrate, but other hard surfaces or mud that can support the weight of the reef are also used (Sellers and Stanley, 1984). A current strong enough to supply food but not re-suspend sediment is necessary for survival (Sellers and Stanley, 1984; EOBRT, 2007). Once, a suitable attachment site is found, the oysters reabsorb the foot and velum and become sedentary spat (Sellers and Stanley, 1984). Settlement can be repeated until suitable habitat is found, but once metamorphosis is complete, the oyster is permanently sessile (EOBRT, 2007). Therefore, adult distribution is determined by the factors effecting larval dispersal, settlement, and survival. Oyster Drills, Channeled Whelk, starfish, crabs, and the Flat Worm, *Stylochus ellipticus*, are predators of small oysters (Sellers and Stanley, 1984). Adult oysters are sessile and usually occur in groups called beds or reefs in a variety of habitats including: oceanic bays, estuaries, drowned river mouths, and behind barrier beaches with wide ranges in temperature, salinity, and water clarity (Sellers and Stanley, 1984). The extent of Eastern Oyster distribution to the north and inland in the northern extent of their range are temperature limited, and predation and food availability also play a role in distribution and survival (EOBRT, 2007). Eastern Oysters are filter feeders of phytoplankton and suspended detritus (Sellers and Stanley, 1984). Oyster Drills and Channeled Whelks are common predators of adults (Sellers and Stanley, 1984). Eastern Oyster is a keystone species in most estuaries throughout their range, providing habitat for many managed and unmanaged species and promoting species diversity, effecting water circulation patterns, and improving water quality (EOBRT, 2007). Harvesting of most natural beds in the northeastern United States is no longer cost-effective (Sellers and Stanley, 1984; EOBRT, 2007). Aquaculture is a major source of commercial oysters (EOBRT, 2007). Eastern Oysters are managed by each state in the United States and by Canadian and Caribbean governing bodies. A petition to list Eastern Oyster as endangered or threatened was denied because the long-term persistence of oyster throughout its range is not believed to be at risk in the foreseeable future (EOBRT, 2007).

Literature Cited:

- Eastern Oyster Biological Review Team (EOBRT). Status review of the eastern oyster (*Crassostrea virginica*). Report to the National Marine Fisheries Service, Northeast Regional Office. 2007. NOAA Tech. Memo. NMFS F/SPO-88, 105 p. Available: [http://www.nmfs.noaa.gov/pr/species/Status%20Reviews/eastern\\_oyster\\_sr\\_2007.pdf](http://www.nmfs.noaa.gov/pr/species/Status%20Reviews/eastern_oyster_sr_2007.pdf)
- Kimmel DG, Newell RI. The influence of climate variation on eastern oyster (*Crassostrea virginica*) juvenile abundance in Chesapeake Bay. *Limnol Oceanogr.* 2007; 52(3): 959-965. doi: 10.4319/lo.2007.52.3.0959
- Kimmel DG, Tarnowski M, Newell RI. Long-term (1939 to 2008) spatial patterns in juvenile eastern oyster (*Crassostrea virginica*, Gmelin 1791) abundance in the Maryland portion of Chesapeake Bay. *J Shell Res.* 2012; 31(4), 1023-1031. doi: <http://dx.doi.org/10.2983/035.031.0414>
- Ries JB, Cohen AL, McCorkle DC. Marine calcifiers exhibit mixed responses to CO<sub>2</sub>-induced ocean acidification. *Geol.* 2009; 37(12), 1131-1134. doi: 10.1130/G30210A.1

Sellers MA, Stanley JG. Species profiles: life histories and environmental requirements of coastal fishes and invertebrates (North Atlantic) – American oyster. 1984. U.S. Fish Wildl. Serv. FWS/OBS-82/11.23. U.S. Army Corps of Engineers, TR EL-82-4. 15 pp. Available: <http://www.nwrc.usgs.gov/publications/specprof.htm>

Talmage SC, Gobler CJ. The effects of elevated carbon dioxide concentrations on the metamorphosis, size, and survival of larval hard clams (*Mercenaria mercenaria*), bay scallops (*Argopecten irradians*), and Eastern oysters (*Crassostrea virginica*). *Limnol Oceanogr.* 2009; 54(6): 2072-2080. doi: 10.4319/lo.2009.54.6.2072

Waldbusser GG, Voigt EP, Bergschneider H, Green MA, Newell RI. Biocalcification in the eastern oyster (*Crassostrea virginica*) in relation to long-term trends in Chesapeake Bay pH. *Estuar Coasts.* 2011; 34(2): 221-231. 10.1007/s12237-010-9307-0