

Synthesis Element 5: Past, Current and Projected Changes in Watershed and Tidal Water Temperatures and Implications for Ecosystem Processes Influencing Stream, River and Estuarine Health

At-a-Glance Summary

- Chesapeake Bay watershed air temperatures and stream-water temperatures have been rising since the 1960s and at higher rates during the 1985-2010 period compared with the 1961-1985 period.
- Stream-water temperatures have been rising faster than air temperatures across the Chesapeake Bay watershed, indicating land use-based factors are also influencing stream-water temperatures.
- Chesapeake Bay tidal water temperatures have been increasing over the past three decades, driven largely by atmospheric forcings and the warming ocean boundary.
- These increasing watershed and tidal water temperatures have significant implications for aquatic living resources and the underlying biological and physical processes which directly influence habitat suitability.

A. Contributors

Rich Batiuk, CoastWise Partners; Nora Jackson, Chesapeake Research Consortium/Chesapeake Bay Program Office; John Clune, United States Geological Survey; Kyle Hinson, Virginia Institute of Marine Science; Renee Karrh, Maryland Department of Natural Resources; Mike Lane, Old Dominion University; Rebecca Murphy, University of Maryland Center for Environmental Science/Chesapeake Bay Program Office; and Roger Stewart, Virginia Department of Environmental Quality.

B. Resources

Published papers cited as references; Chesapeake Bay water quality monitoring network's long term trend analyses generated by Rebecca Murphy, Renee Karrh, and Mike Lane; U.S. Environmental Protection Agency Climate Change Indicator development documentation; Pennsylvania Report on Climate Impacts; and interviews with recognized regional scientists and data analysts.

C. Approach

Synthesized evidence for long term changes in watershed and tidal Bay water temperatures, then engaged researchers and statistical analysts currently involved in in-depth analysis and evaluation of

both the trends and the likely underlining causes behind the observed trends and finished with accounting for the implications for the watershed and estuarine ecosystem.

D. Synthesis



Watershed and Tidal Bay Water Temperature Trends

Watershed and tidal Bay water temperatures are rising and have been for the past several decades. Preston (2004) reported an average Bay water temperature increase of $\sim 0.8\text{-}1.1^\circ\text{C}$ from 1949-2002 as derived from direct observations and satellite measurements. Ding and Elmore (2015) found increases in Chesapeake Bay surface water temperature of $\sim 0.4\text{-}2^\circ\text{C}$ from 1984-2010, also based on direct observations and satellite measurements. U.S. Geological Survey trend analysts reported that average non-tidal stream temperatures increased 2.52°F from 1960 to 2010, while air temperatures increased 1.99°F (Rice and Jastram 2015).

Non-Tidal Water Temperature Trends

Key takeaways from trend analysis of monthly mean air temperature at 85 sites and instantaneous stream-water temperature at 129 sites within or near the Chesapeake Bay watershed from 1960 to 2010 (Rice and Jastram 2015) (Figure V-4) include:

- Analysis of both air and stream-water temperatures for two periods, 1961–1985 and 1985–2010, relative to the climate normal period of 1971–2000, indicate that the 1985-2010 period was statistically significantly warmer than the 1961-1985 period for both mean air temperature and stream-water temperature;
- Across the Chesapeake Bay watershed and the surrounding region, statistically significant temporal trends of 0.023°C per year for air temperature and 0.028°C per year for stream-water temperature were determined;
- From 1960 through 2010, water temperature increased significantly at 53 of 129 stations analyzed in the region;
- Stream-water temperature decreased significantly at 7 of those 129 stations over the same period;
- In areas where major dams were and the land cover was principally deciduous forest, stream-water temperatures were increasing slower than air temperatures, whereas agriculture-dominated regions in the absence of major dams were correlated with stream-water increasing faster than air temperatures;
- Increasing stream-water temperature trends are detected despite increasing trends in streamflow in the northern Chesapeake Bay watershed and surrounding region; and
- Increases in water temperature occurred at the greatest rates in the southern Chesapeake Bay watershed and surrounding region.



Filled shapes represent statistically significant trends.
 Open shapes represent trends that are not statistically significant.

Figure V-1. Changes in Stream Water Temperatures in the Chesapeake Bay Region, 1960–2010.

Source: Rice and Jastram 2015


The map in Figure V-1 shows the change in water temperature at 129 stream gauges across the Chesapeake Bay region from 1960 to 2010. Red circles show locations where temperatures have increased; blue circles show locations where temperatures have decreased (Rice and Jastram 2015). Filled circles represent sites where the change was statistically significant based on the U.S. EPA Climate Indicator¹.

Water temperature in streams can be affected by factors other than climate, including industrial thermal discharges, hydrologic alteration (for example, channelization, piping, and impoundment), land cover, location, and topography. A more detailed analysis of this data set found that water temperature tends to increase more quickly than air temperature in agricultural areas without major dams, but more slowly at forested sites and in areas influenced by dams (Rice and Jastram, 2015). For this indicator, water temperature measurements from all available stream gages with appropriate records within the study area were used, as described in Rice and Jastram (2015), regardless of potential influences from anthropogenic disturbances.

A comparison, using the Rank-Sum test (Helsel and Hirsch, 2002), of relatively undisturbed reference stations (n = 35), as determined by Falcone (2011), with all other stations (n = 94) in the dataset demonstrated no significant difference (alpha = 0.05) in trends between the two groups of stations. Trends were determined using ordinary least-squares linear regression of sites-specific monthly water temperature anomalies, as described by Rice and Jastram (2015). The Cochrane-Orcutt method (Cochrane and Orcutt, 1949) was used to remove the effect of serial correlation, thus allowing determination of the statistical significance of water temperature trends at individual stations. Of the 129 stations analyzed, 60 (47 percent) had trends that were significant to a 95-percent level ($p \leq 0.05$), including 53 stations with temperature increases and seven with decreases.

Sources of variability include localized factors such as topography, geology, elevation, and natural land cover within individual watersheds. Variability between individual temperature measurements could result from variations in weather—for example, if a recent storm led to an increase in streamflow. Additionally, some sites may be more affected by direct human influences (such as land-cover and land use change or hydrologic modification) than others and does not include any sites that are affected by tides.

The Virginia Department of Environmental Quality operates a network of 410 permanent trend stations where monthly or bimonthly data are collected for a variety of key water quality parameters. These fixed stations are located in areas of special interest including those near the mouths of our major rivers, along the fall line, near flow gaging stations, at designated non-tidal stations monitored to evaluate how rivers affect the Chesapeake Bay. In the 2018 Integrated Report on Water Quality Trends in Virginia from 1997-2016, water temperature was included in the trend analysis as a water quality indicator variable (Figure V-2) (Steward 2018). Temperature has an influence on regulating respiration rates, spawning, and the maximum concentration of dissolved oxygen in solution with the ambient



water (increasing temperature reduces dissolved oxygen saturation in water and, therefore, may limit respiration). In addition, animals and plants under thermal stress from high-water temperatures are at increased risk of adverse effects from other pollutants. Temperature standards exist for “the propagation and growth of a balanced indigenous population of aquatic life” as described in the Clean Water Act (33 U.S.C. §1251 et seq; this is, more correctly, a balanced and indigenous community of aquatic life). Pollution events that cause harm to aquatic communities via water cooling are extremely rare in VA, and not known to exist at the stations in the trend network. Therefore, increasing trends in water temperature are considered degradation, and decreases in temperature are considered improvements (Stewart 2018).




Figure V-1. Observed annual averaged surface and bottom water temperatures across Chesapeake Bay from 1985 through 2020. Source: Hinson et al. 2021.

Tidal Bay Water Temperature Trends

Using estimates of changes from downscaled global climate models (GCMs) and the Chesapeake Bay Program Partnership’s modeling framework, Tian et al. (2021) documented and projected changes in Chesapeake Bay water temperatures of 0.85-0.9°C from 1995-2025. When Hinson et al. (2021) used a






Figure V-3. Observed annual averaged surface and bottom water temperatures across Chesapeake Bay from 1985 through 2020. Source: Hinson et al. 2021.

combination of observations and model outputs to report that throughout Chesapeake Bay’s mainstem, similar warming rates were found at the surface and bottom between the late 1980s and late 2010s of 0.02 °C per year, with elevated summer rates (0.04 °C per year) and lower rates of winter warming (0.01 °C per year) (Figure V-3). These annual rates yielded an annual average Bay-wide warming of ~0.7°C throughout the Chesapeake Bay’s water column over the past 30-year period, with a 1.0 °C increase during the summertime and a 0.3°C during the winter months over the same three-decade period (Hinson et al. 2021).

Recent work by Murphy and colleagues (personal communication), using generalized additive model approach to evaluating water quality as described in Murphy et al. 2019, yielded the 1985-2019 estimated changes in tidal water bottom and surface temperatures seen in Figures V-4 and V-5, respectively.



Driving Forces Behind Warming of Chesapeake Bay Tidal Waters

Hinson et al. (2021) have identified four principal mechanisms responsible for the observed increasing

Figure V-4. Long term flow-adjusted trends in surface water temperatures at the Chesapeake Bay Mainstem and Tidal Tributary Water Quality Monitoring Program stations through 2019.

Figure V-5. Long term flow-adjusted trends in bottom water temperatures at the Chesapeake Bay Mainstem and Tidal Tributary Water Quality Monitoring Program stations through 2019.

temperatures of Chesapeake Bay’s tidal waters, listed here in the order of their relative influence: atmospheric forcings, warming ocean boundary, sea level rise and increasing river temperatures (Figure V-6).

Hinson et al. 2021 utilized “the extensive observational network of in situ data along with a watershed-estuarine modeling system forced by realistic atmospheric and oceanic inputs to quantify and better understand the causes of warming in the Chesapeake Bay. Using this approach, a more robust estimate of the recent observed temperature trends and the causality of said trends can be more precisely determined.”





Figure V-6. Illustration of the four major mechanisms driving changes in water temperature throughout the Chesapeake Bay's mainstem, tidal tributaries and embayments. Source: Hinson et al. 2021

Temperature changes were largely very similar at the Bay's and tidal tributaries' surface and bottom of the water column (Hinson et al. 2021) (Figure V-7). Some regional differences in temperature changes were reported, with higher temperature changes estimated for the Susquehanna Flats and adjoining upper Bay mainstem, the lower Bay and mouth of the Bay, and the tidal fresh reaches of the major tidal tributaries (Figure V-8). There is evidence supporting river temperature influences in the upper tidal fresh reach of the major tidal tributaries and the upper Chesapeake Bay—Susquehanna Flats and the upper Bay mainstem reach down to about Back River on the western shore (see Figure V-7).




There is substantial variation in the estimated water column temperature changes over the past 30 years between months, with generally more warming of water temperatures from May-October than November-April (Figure V-9). The observed increasing river temperatures are estimated produce very limited warming of water column temperatures in the Bay's mainstem (Hinson et al. 2021) (Figure V-10).

Figure V-7. Two-dimensional depth profile of the 30-year change in water column temperature along the Chesapeake Bay mainstem.

Sea level rise is estimated to slightly cool Bay mainstem water column temperatures from April through September, and result in the warming of bottom Bay mainstem waters in the winter months (November through February) (Figure V-10) (Hinson et al. 2021). Increasing ocean temperatures are estimated to contribute significantly to the summer warming of the Bay water column temperatures between June and October, with a small effect on water column temperature for the remaining months of the year (Figure V-10). Atmospheric forcings are estimated to play biggest role in driving increasing water column temperatures throughout the Bay’s tidal waters, but the effects on water temperatures are lessened during summer months of July through September and contribute to a cooling of Bay water temperatures during December (Figure V-10).

Figure V-8. The 30-year change in observed water temperatures at the surface and bottom across Chesapeake Bay. Source: Hinson et al. 2021.







Figure V-10. Model-simulated 30-year bottom water temperature change throughout Chesapeake Bay by month compared with model-simulated bottom water temperature change estimated to be caused by river temperatures, sea level rise, ocean temperature and atmospheric forcings. Source: Hinson et al. 2021.

Atmospheric warming is the dominate influence on increasing Bay water column temperature almost everywhere across the tidal waters, contributing about 78% to the combined effect on changes in bottom Chesapeake Bay water temperatures observed by the past 30 years, equal to about a 0.6°C change over this timeframe (Figure V-11).

The warming the adjacent Atlantic Ocean plays a large role in the changes in southern Chesapeake Bay’s water temperatures, with about a 26% contribution to the overall Bay bottom temperatures over the past three decades. Ocean warming alone has contributed at least 50% or greater to the increased Bay water column temperature during the summer months over the past 30 years (Hinson et al. 2021). The increasing temperatures in the rivers flowing into Chesapeake Bay only influence the water column temperatures of the immediate tidal fresh reaches of the tidal tributaries, making no measurable contribution to observed changes in bottom Chesapeake Bay water temperatures observed by the past 30 years. Sea level rise is estimated to slightly cool Bay water column temperatures across the tidal waters, contributing an offsetting 6% cooling contribution to the overall Bay bottom temperatures over the past three decades, about 0.1°C difference over this timeframe (Hinson et al. 2021).



Implications for Ecosystem Processes

Watershed Ecosystem Processes

Water temperature affects all chemical and biological processes of aquatic organisms, as well as being directly linked to survival for temperature-sensitive organisms like brook trout. Water temperature integrates what is happening on the land (e.g. forested, urban impervious), and affects the way nutrients and other pollutants behave in the water column.

Temperatures can vary naturally along the length of a stream, from cold temperatures near a source of meltwater to higher temperatures near its outlet to the tidal water. The temperature at any given point is a product of many different factors, including sources of water (for example, melted snow, a recent rainstorm, or groundwater), the amount of water in the stream (streamflow), air temperature, plants along the bank (for example, trees that provide shade), and the amount of development within the watershed. Over time, however, an area's climate has the strongest natural influence on a stream's temperature. Higher temperatures reduce levels of dissolved oxygen in the water, which can negatively affect the growth and productivity of aquatic life. Persistently warmer temperatures in streams can accelerate natural chemical reactions and release excess nutrients into the water.

Despite the wide variability of the streams and rivers with respect to watershed area, channel geometry, aspect, elevation, thermal capacity, the presence or absence of riparian buffers, microclimate conditions, and land cover, on the whole, water temperature increased from 1960 to 2010. For sites with significantly increased water temperature, 85 % of the variability could be explained by increased atmospheric temperature, despite increased streamflow at some sites (Rice and Jastram 2015).

Estuarine Ecosystem Processes

Tian et al. 2021 reported that increasing Chesapeake Bay water column temperatures will result in reduced oxygen saturation, increased biological rates, and increased stratification of the water column. Their research focused on better understanding how changing oxygen solubility affects dissolved oxygen concentrations in the bottom waters of a stratified Chesapeake Bay.

Higher water temperature reduce the amount of oxygen which can become soluble in water, forming dissolved oxygen. The higher water temperatures will also increase the remineralization rate, that is the natural bacterial decomposition of organic matter into nitrogen, phosphorus and carbon, internally fueling growth of algae. Both of these processes lead to further expansion of and sustaining existing hypoxic (low dissolved oxygen) and anoxic (no dissolved oxygen) conditions in the deeper bottom waters of the Bay mainstem and lower tidal tributaries.

Running a series of scenario simulations using Chesapeake Bay Water Quality Model to determine the magnitude of various mechanisms controlling the effect of increasing water temperature on dissolved oxygen in the Chesapeake Bay, Tian et al. 2021 reported the following findings. They estimated the average hypoxic volume in the summer would increase by 9% from 1995 to 2025 as air temperature increases by 1.06°C and water temperature by 0.9°C. Of the three major drivers of water temperature change impacts, the change in dissolved oxygen solubility contributes 55% to the model projected change in hypoxic volume, biological rates 33%, and stratification 11%.

Off the mouth of the Rappahannock River, the abrupt change in bathymetry and “the convergence between seaward-moving freshwater and landward-moving saltwater causes downwelling and enhanced vertical mixing which introduces surface water of higher temperature to the deep channel and accelerates organic matter remineralization and oxygen consumption in deep waters” (Tian et al. 2021). As surface water dissolved oxygen concentrations will decrease under continued warming of the climate due to lower oxygen solubility, surface waters with even lower dissolved oxygen concentrations will flux to the deep channel further exacerbating development of low to no dissolved oxygen conditions in the deep channel of Chesapeake Bay.

Hinson et al. 2021 reported “on average during the period from May to October, a time of particular interest since it encompasses the bottom hypoxia season, there is more warming in the shallower southernmost extent of the mainstem than in the rest of Chesapeake Bay. Combined with the findings reported by Tian et al. 2021, the warming of the southern Chesapeake Bay mainstem waters will further exacerbate the increased impact of warming water temperatures on low and no dissolved oxygen conditions in the deeper channels of the Chesapeake Bay mainstem to the north.

E. Evaluation

Key Findings

A U.S. Geological Survey analysis of non-tidal stream temperatures from 1960 through 2010 documented that average non-tidal stream temperatures increased 2.52 °F from 1960 to 2010, while air temperatures increased 1.99 °F (Rice and Jastram 2015). These major findings were:

- Analysis of both air and stream-water temperatures for two periods, 1961–1985 and 1985–2010, relative to the climate normal period of 1971–2000, indicate that the 1985-2010 period was statistically significantly warmer than the 1961-1985 period for both mean air temperature and stream-water temperature;
- Across the Chesapeake Bay watershed and the surrounding region, statistically significant temporal trends of 0.023 °C per year for air temperature and 0.028 °C per year for stream-water temperature were determined;
- From 1960 through 2010, water temperature increased significantly at 53 of 129 stations analyzed in the region;
- Stream-water temperature decreased significantly at 7 of those 129 stations over the same period;
- In areas where major dams were and the land cover was principally deciduous forest, stream-water temperatures were increasing slower than air temperatures, whereas agriculture-dominated regions in the absence of major dams were correlated with stream-water increasing faster than air temperatures;
- Increasing stream-water temperature trends are detected despite increasing trends in streamflow in the northern Chesapeake Bay watershed and surrounding region; and
- Increases in water temperature occurred at the greatest rates in the southern Chesapeake Bay watershed and surrounding region.

Rice and Jastram (2015) concluded “continued warming of contributing streams to Chesapeake Bay likely will result in shifts in distributions of aquatic biota and contribute to worsened eutrophic conditions in the bay and its estuaries.”

There is significant evidence of widespread increases in Chesapeake Bay water column temperatures reported independently by an array of different research and data analysis teams over the past decade. And recently, a research team composed of scientists from the Virginia Institute of Marine Science and Penn State University published an in-depth evaluation of the major drivers for the observed increases in Chesapeake Bay water column temperatures (Hinson et al. 2021). Their major findings are summarized as:

- Atmospheric forcings and warming ocean boundary are the most pertinent driving forces to future warming of Chesapeake Bay water temperatures;
- Atmospheric forcings (air temperature increases/decreases) main driver influencing Bay water temps year-round, but effects lessened during summer;
- Warming ocean boundary effects are important in summer (influenced => 50% warming), but small otherwise during the rest of seasons;
- Sea level rise slightly cools Chesapeake Bay mainstem waters from April-September and warms bottom waters in winter;

- River temperatures produce little to no warming in the Chesapeake Bay's mainstem, but still influence temperature in the tidal fresh and low salinity waters in the upper reaches of the tidal tributaries and embayments; and
- Future warming of Chesapeake Bay waters will depend not only on global temperature trends, but also on regional circulation patterns in mid-Atlantic waters which are currently warming faster than the atmosphere.

Tian et al. 2021 warned that increasing Chesapeake Bay water temperatures will result in increased volumes of low dissolved oxygen due to direct effects on oxygen solubility, biological processes rates and water column stratification.

Management Implications

For freshwaters, there are implications for potential shifts in floral and faunal species distributions. Streams at the upper end of the water temperature distribution may become unsuitable habitat for certain cool-water fish species (Eaton and Scheller 1996; Isaak et al. 2012). Increasing water temperature also may make some streams suitable for species not currently present, allowing warm-water species, including invasive species and pathogens, to move into previously cool-water habitats. Streams draining forested watersheds with major dams warmed more slowly than other watersheds and are likely to become even more important as refugia for cool-water species in a warming world (Rice and Jastram 2015). In addition, warmer water temperatures in the watershed's streams and rivers could decrease the availability of water used for power plant cooling and could have other interactions with built infrastructure².

Reducing the water temperatures of the river flowing into Chesapeake Bay will have no to a very minimal to affect the continued warming of most of Chesapeake Bay's water column temperatures. River water temperatures do influence the water temperatures of the tidal tributary reaches just down tide of the river inputs as well as the Susquehanna Flats and the upper Bay mainstem reach down to about Back River on the western shore. These tidal fresh reaches provide for important spawning, nursery and year-round habitats for anadromous (e.g., striped bass), semi-anadromous (e.g., white perch) and resident (e.g., largemouth bass) fish populations which are directly affected by changes in tidal water temperature.

Changing the magnitude of the two major influences on Bay water temperatures—atmospheric forcings and ocean warming—are clearly management and human behavioral challenges to be addressed at the global to local scales, collectively. However, the resultant effect of warmer water temperatures on biological, chemical and other ecosystem process will very likely require additional nutrient and sediment load reductions to mitigate these impacts on the Bay's living resources. Continue warming of the Bay waters will affect the temperature thresholds critical to the survival, growth, behavior and migration patterns of individual species and entire communities as well as their prey and the suitability of their surrounding habitats. Further research on the warming of Chesapeake

² Pennsylvania Climate Impacts Assessment.

Bay waters must not only better understand the impacts on water temperature from atmospheric changes, but also changes in the adjacent coastal ocean.

F. Bibliography

References Cited

Cochrane, D. and G.H. Orcutt. 1949. Application of Least Squares Regression to Relationships Containing Auto-Correlated Error Terms. *Journal of the American Statistical Association*, 44, 32-61.

Ding, H. and A.J. Elmore. 2015. "Spatio-Temporal Patterns in Water Surface Temperature from Landsat Time Series Data in the Chesapeake Bay, U.S.A." *Remote Sensing of Environment* 168: 335–348. <https://doi.org/10.1016/j.rse.2015.07.009>.

Eaton and Scheller. 1996. "Effects of climate warming on fish thermal habitat in streams of the United States" *Limnology and Oceanography* 41 (5), 1109-1115. <https://doi.org/10.4319/lo.1996.41.5.1109>

Falcone. 2011. "GAGES-II: Geospatial Attributes of Gages for Evaluating Streamflow" https://water.usgs.gov/lookup/getspatial?gagesII_Sept2011

Helsel, D.R., Hirsch, R.M., Ryberg, K.R., Archfield, S.A., and Gilroy, E.J. 2020. Statistical methods in water resources: U.S. Geological Survey Techniques and Methods, book 4, chapter A3, 458 p., <https://doi.org/10.3133/tm4a3>. [Supersedes USGS Techniques of Water-Resources Investigations, book 4, chapter A3, version 1.1.]

Hinson, K., M.A.M. Friedrichs, P. St-Laurent, F. Da, R.G. Najjar. 2021. "Extent and Causes of Chesapeake Bay Warming." *Journal of the American Water Resources Association* 1–21. <http://doi.org/10.1111/1752-1688.12916>

Isaak, D.J., Wollrab, S., Horan, D. *et al.* Climate change effects on stream and river temperatures across the northwest U.S. from 1980–2009 and implications for salmonid fishes. *Climatic Change* 113, 499–524 (2012). <https://doi.org/10.1007/s10584-011-0326-z>

Jastram, J.D., and K.C. Rice. 2015. Air- and stream-water-temperature trends in the Chesapeake Bay region, 1960–2014: U.S. Geological Survey Open-File Report 2015–1207, 28 <http://dx.doi.org/10.3133/ofr20151207>.

Linker, L. C., 2020. "Influence of Climate Change Risk on the Chesapeake Bay Open-Water Dissolved Oxygen Water Quality Standard." For distribution to the Chesapeake Bay Program Modeling Workgroup and Water Quality Goal Implementation Team. July 29, 2020. Chesapeake Bay Program Office, Annapolis, MD.

Murphy, R.R., E. Perry, J. Harcum, and J. Keisman. 2019. A Generalized Additive Model Approach to Evaluating Water Quality: Chesapeake Bay Case Study. *Environ. Modelling Software* 118: 1-13. <https://doi.org/10.1016/j.envsoft.2019.03.027>.

Preston, B.L. 2004. Observed Winter Warming of the Chesapeake Bay Estuary (1949-2002): Implications for Ecosystem Management. *Environmental Management* 34: 125–139.

<https://doi.org/10.1007/s00267-004-0159-x>.

Rice, K.C., and J.D. Jastram. 2015. Rising air and stream-water temperatures in Chesapeake Bay region, USA: *Climatic Change*, v. 128, p. 127-138. <https://doi.org/10.1007/s10584-014-1295-9>

Stewart, R. 2018. *Integrated Report on Water Quality Trends in Virginia*. Virginia Department of Environmental Quality, Richmond, Virginia.

Tian, R., C.F. Cerco, G. Bhatt, L.C. Linker, and G.W. Shenk. 2021. Mechanisms Controlling Climate Warming Impact on the Occurrence of Hypoxia in Chesapeake Bay. *Journal of the American Water Resources Association* 1–21. <https://doi.org/10.1111/1752-1688.12907>.

Additional Resources

Kyle Hinson’s presentation on the “Extent and Causes of Chesapeake Bay Warming” as presented to the Chesapeake Bay Program Partnership’s Modeling Workgroup can be accessed at:

https://www.chesapeakebay.net/channel_files/42529/hinson_bay_warming_-_20210407.pdf

The “baytrendsmap” link that can be used to generate custom maps and explore the GAM trend analysis results is accessible at: <https://baytrends.chesapeakebay.net/baytrendsmap/>

EPA Climate change indicator can be accessed at:

<https://www.epa.gov/climate-indicators/climate-change-indicators-stream-temperature>

Pennsylvania Climate Impacts Report can be accessed at:

https://files.dep.state.pa.us/Energy/Office%20of%20Energy%20and%20Technology/OETDPortalFiles/Climate%20Change%20Advisory%20Committee/2020/12-22-20/2021_IA_Draft_Final_12-15-20.pdf

Stream temperature EPA technical documentation can be accessed at:

https://www.epa.gov/sites/production/files/2016-08/documents/stream-temperature_documentation.pdf