

Machine learning and differentiable modeling for Geosciences

Chaopeng Shen

¹Civil and Environmental Engineering
Penn State University
cshen@engr.psu.edu



<https://github.com/mhpi>
Hydroml.org

Hydroml.org
HydroML Symposium, May 22-26, 2022, Penn State
HydroML 2, May 2023, Berkeley, CA

About me

- Ph.D. Michigan State in Env. Engr.
- Postdoc Lawrence Berkeley National Lab
- Associate Editor, Water Resources Research
Specialty Chief Editor, Frontiers in Water:
Water and AI.
- “Grew up” as a process-based modeler,
solving PDEs. See both sides of the story.
- Got into ML since 2016.




Overview

- **What** ML models have we got comfortable with?
- **What** is the fundamental strengths of ML models compared to process-based models?
- **What** is *differentiable modeling (DM) in geosciences*?
- **What** can DM bring into global hydrology?

nature reviews earth & environment

<https://doi.org/10.1038/s43017-023-00450-9>

Perspective


 Check for updates

Differentiable modelling to unify machine learning and physical models for geosciences


Shen et al., 2023 Nature Reviews Earth & Environment <https://t.co/qyuAzYPA6Y>

A list of authors and their affiliations appears at the end of the paper

Process-based Earth-system models were highly valuable but some challenges emerged...



- Increasing complexity
- Difficult to evolve quickly with more big data.
- May contain problematic assumptions.
- Influenced by human intuition & biases



What is DL and why DL?

a rebranding of neural networks featuring

- (i) Large capacity
- (ii) Hidden layers that automatically extract features
- (iii) Improved architecture/regularization
- (iv) Working directly with data

a primary value proposition is the avoidance of expertise!

Three phases

1. Use ML to learn where the limit is.
2. Understand the gaps in our knowledge.
3. Using ML to unify across domains.



Water Resources Research

AN AGU JOURNAL


Review Article | [Open Access](#)

A trans-disciplinary review of deep learning research and its relevance for water resources scientists


Chaopeng Shen [✉](#)

First published: 30 August 2018 | <https://doi.org/10.1029/2018WR022643>

$X \rightarrow Y$




Some basic deep learning architectures



Hydrologic DL phase 1.

A hydrologic model w/o structural assumptions...



→ Free from structural assumptions


→ A chance to start anew!

→ A chance to see where the limit is!




Case studies— first phase of DL in water


- Soil Moisture Active Passive (SMAP)
 - Launched recently (2015/04)
 - 2~3 days revisit time
 - Senses moisture-dependent top surface soil
- Streamflow modeling
 - Daily data
 - Accompanying attributes
 - With reservoirs, in data-sparse regions
- Dissolved oxygen
- Water temperature
- Sediment
- Snow water equivalent



Gridded models



Basin-scale models



Long-term projections (first-phase of DL in hydrology)

- Examined comparison with in-situ data & long-term projections


Geophysical Research Letters

Research Letter | Full Access

Prolongation of SMAP to Spatiotemporally Seamless Coverage of Continental U.S. Using a Deep Learning Neural Network

Kuai Fang, Chaopeng Shen, Daniel Kifer, Xiao Yang

First published: 16 October 2017 | <https://doi.org/10.1002/2017GL075619> | Cited by: 3



Fang et al., 2017

Water Resources Research


RESEARCH ARTICLE
10.1029/2019WR026793

Special Section:
Big Data & Machine Learning
in Water Sciences: Recent
Progress and Their Use in
Advancing Science

Enhancing Streamflow Forecast and Extracting Insights Using Long-Short Term Memory Networks With Data Integration at Continental Scales


Dapeng Feng¹, Kuai Fang^{1,2}, and Chaopeng Shen¹

¹Civil and Environmental Engineering, Pennsylvania State University, State College, PA, USA, ²Now at: Earth System Science, Stanford University, Stanford, CA, USA




Short-term forecast





- Traditional “data assimilation” scheme




Simulation \rightarrow Observation – (ENKF) \rightarrow Correction

Choices: covariance matrix, what to include, how to solve, bias correction, etc.





Journal of Hydrology
 Available online 16 May 2021, 126455
 In Press, Journal Pre-proof



Research papers

Continental-scale streamflow modeling of basins with reservoirs: towards a coherent deep-learning-based strategy


Wenyu Ouyang^a, Kathryn Lawson^b, Dapeng Feng^b, Lei Ye^a, Chi Zhang^a, Chaopeng Shen^b


Sparse-data region

- Transfer learning

RESEARCH ARTICLE

How to enhance hydrological predictions in hydrologically distinct watersheds of the Indian subcontinent?

Nikunj K. Mangunkiya¹ | Ashutosh Sharma¹  | Chaopeng Shen²



Ma et al., WRR

<https://doi.org/10.1029/2020WR028600>

Basis of








FIGURE 1 Locations of the gauge station and watershed across India. The gauges are spread across different Koppen climate zones and have distinct watershed characteristics.

Multiscale soil moisture – learning from two teachers



Test period: 2015-04-01 to 2020-03-31



Geophysical Research Letters®

Research Letter | Full Access

A multiscale deep learning model for soil moisture integrating satellite and in-situ data

Jiangtao Liu, Farshid Rahmani, Kathryn Lawson, Chaopeng Shen

First published: 14 March 2022 | <https://doi.org/10.1029/2021GL096847>

Water quality

nature water

Explore content ▾ About the journal ▾ Publish with us ▾ Subscribe

[nature](#) > [nature water](#) > [articles](#) > article


Article | Published: 09 March 2023

Temperature outweighs light and flow as the predominant driver of dissolved oxygen in US rivers

Wei Zhi, Wenyu Ouyang, Chaopeng Shen & Li Li

Nature Water 1, 249–260 (2023) | [Cite this article](#)

Dissolved Oxygen



ELSEVIER

Contents lists available at [ScienceDirect](#)

Science of the Total Environment

journal homepage: www.elsevier.com/locate/scitotenv



A deep learning-based novel approach to generate continuous daily stream nitrate concentration for nitrate data-sparse watersheds

Gourab Kumer Saha^a, Farshid Rahmani^b, Chaopeng Shen^b, Li Li^b, Raj Cibin^{a,b,*}

^a Department of Agricultural and Biological Engineering, The Pennsylvania State University, United States of America

^b Department of Civil and Environmental Engineering, The Pennsylvania State University, United States of America

Nitrate

ENVIRONMENTAL RESEARCH LETTERS

LETTER

Exploring the exceptional performance of a deep learning stream temperature model and the value of streamflow data

Farshid Rahmani¹, Kathryn Lawson¹, Wenyu Ouyang², Alison Appling³, Samantha Oliver⁴ and Chaopeng Shen¹


¹ Civil and Environmental Engineering, Pennsylvania State University, University Park, State College, PA, United States of America

² School of Hydraulic Engineering, Dalian University of Technology, Dalian, People's Republic of China

³ US Geological Survey, Reston, VA, United States of America

⁴ US Geological Survey, Upper Midwest Water Science Center, Middleton, WI, United States of America

Water temperature




pure DL models



Human modelers

We need hybrids



Phase 2: How to surpass the teacher (training data)

Training data often have limitations:

Resolution, accuracy, time interval, availability (unobserved variables), geographical imbalance, not enough extremes, not capturing nonstationarity...

How to overcome such limitations?

- Inclusion of physics
- Learning about physics.



Similarity & Differences between deep learning (DL) and process-based models (PBM)?



[This Photo](#) by Unknown Author is licensed under [CC BY-NC](#)

Purely data-driven NNs	Purely process-based models
Similarities	
$y = g^W(u, x, A)$ $W = \operatorname{argmin}(L(y, y^*))$	$y = f^\theta(u, x, A)$ $\theta = \operatorname{argmin}(L(y, y^*))$




[This Photo](#) by Unknown Author is licensed under [CC BY-SA](#)

The secret? Differentiable programming!

What does “Differentiable” mean?

- The ability to rapidly compute gradients $\frac{dL}{d\theta}$
- Enabling training by gradient descent

Automatic differentiation



$$e = a * b + c \log(d)$$


$$\frac{\partial e}{\partial a} = b(1) = b = 3$$

$$\frac{\partial e}{\partial b} = a(1) = a = 2$$


$$\frac{\partial e}{\partial c} = \log d \times 1 = \log d = 2.3$$

$$\frac{\partial e}{\partial d} = \frac{1}{d} \times c \times 1 = \frac{c}{d} = 0.5$$

Adjoint State method



Differentiable parameter learning



ARTICLE

<https://doi.org/10.1038/s41467-021-26107-z>


OPEN




From calibration to parameter learning: Harnessing the scaling effects of big data in geoscientific modeling

Wen-Ping Tsai¹, Dapeng Feng¹, Ming Pan^{2,3}, Hylke Beck⁴, Kathryn Lawson^{1,5}, Yuan Yang^{6,7}, Jiangtao Liu¹ & Chaopeng Shen^{1,5}✉



(a) PBM or PBM's surrogate (optional)





(b) dPL g_A framework (if historical observations are unavailable)



(a) dPL g_z INFILT




(b) SCE INFILT




Point #1. Data scaling relationships (network effect?)

1. dPL = SCEUA for lowest RMSE
2. dPL scales better with more data
3. Orders of magnitude more efficient
4. (not shown) better results for **untrained** variables and better **spatial generalization** than traditional approach!


Relies on differentiable programming!



What is Differentiable Modeling (DM) in Geosciences?



- NNs mixed w/ process-based equations (priors)
- The priors constrain the learning to an interpretable scope.
- intermediate physical variables.
- Update our knowledge and learn unrecognized relationships from data.



Differentiable, learnable models to learn functions

Hydrol. Earth Syst. Sci., 27, 2357–2373, 2023
<https://doi.org/10.5194/hess-27-2357-2023>
 © Author(s) 2023. This work is distributed under the Creative Commons Attribution 4.0 License.



The suitability of differentiable, physics-informed machine learning hydrologic models for ungauged regions and climate change impact assessment

Dapeng Feng¹, Hylke Beck², Kathryn Lawson¹, and Chaopeng Shen¹

¹Civil and Environmental Engineering, The Pennsylvania State University, University Park, PA, USA

²Physical Science and Engineering, King Abdullah University of Science and Technology, Thuwal, Saudi Arabia

Correspondence: Chaopeng Shen (cshen@engr.psu.edu)


Water Resources Research

Research Article | [Full Access](#)

Differentiable, learnable, regionalized process-based models with multiphysical outputs can approach state-of-the-art hydrologic prediction accuracy

Dapeng Feng, Jiangtao Liu, Kathryn Lawson, Chaopeng Shen

First published: 19 September 2022 | <https://doi.org/10.1029/2022WR032404>





* Not all parameters and detailed processes of HBV sketched here for the sake of simplicity.

Rewritten in PyTorch

Evolve model structure


Approaching LSTM!
But....

- Output untrained variables.
- Multivariate constraints.
- It can help us answer questions!





--- MPR+mHM
 B=492 NSE₅₀: 0.528
 - - - dPL+HBV (δ_1)
 B=492 NSE₅₀: 0.618
 - - - dPL+HBV (δ_1)
 B=671 NSE₅₀: 0.64
 — dPL+evolved HBV (δ_n)
 B=671 NSE₅₀: 0.714
 — dPL+evolved HBV with DP ($\delta_n(\beta^t, \gamma^t)$)
 B=671 NSE₅₀: 0.732
 — LSTM
 B=671 NSE₅₀: 0.748

Baseflow



Evapotranspiration



Caveat: not using the ensemble
-- first iteration. Priors do matter.



PBM



PBM+dPL

Differentiable, Learnable, Regionalized Process-Based Models With Multiphysical Outputs can Approach State-Of-The-Art Hydrologic Prediction Accuracy

Dapeng Feng, Jiangtao Liu, Kathryn Lawson, Chaopeng Shen [✉](#)

First published: 19 September 2022 | <https://doi.org/10.1029/2022WR032404> | Citations: 18

<https://doi.org/10.5194/hess-27-2357-2023>
 © Author(s) 2023. This work is distributed under the Creative Commons Attribution 4.0 License.


Article Assets Peer review Metrics Related articles

Research article | [CC](#) [f](#)

30 Jun 2023

The suitability of differentiable, physics-informed machine learning hydrologic models for ungauged regions and climate change impact assessment

Dapeng Feng, Hylke Beck, Kathryn Lawson, and Chaopeng Shen [✉](#)



<https://doi.org/10.5194/gmd-2023-190>
 © Author(s) 2023. This work is distributed under the Creative Commons Attribution 4.0 License.

Abstract Discussion Metrics

Submitted as: model evaluation paper | [CC](#) [f](#)


05 Oct 2023

Status: this preprint is currently under review for the journal GMD.

Deep Dive into Global Hydrologic Simulations: Harnessing the Power of Deep Learning and Physics-informed Differentiable Models (δ HBV-globe1.0-hydroDL)

Dapeng Feng, Hylke Beck, Jens de Bruijn, Reetik Kumar Sahu, Yusuke Satoh, Yoshihide Wada, Jiangtao Liu, Ming Pan, Kathryn Lawson, and Chaopeng Shen [✉](#)

New model





Model	Median NSE	Median KGE	Median absolute (non-absolute) FLV (%)	Median absolute (non-absolute) FHV (%)	Median low flow RMSE (mm/day)	Median peak flow RMSE (mm/day)	Baseflow index spatial correlation	Median NSE of temporal ET simulation
LSTM	0.73	0.77	40.59 (29.70)	13.46 (-4.19)	0.055	2.56	-	-
SAC-SMA	0.66	0.73	59.40 (46.96)	17.55 (-9.79)	0.081	3.19	-	-
HBV	0.73	0.73	56.53 (50.93)	15.29 (-8.89)	0.074	2.56	0.76	0.59
The mysterious model	0.72	0.75	43.29 (37.61)	13.25 (-4.33)	0.048	2.47	0.83	0.61


What can DM bring to global hydrology?

- Spatial extrapolation in data-sparse regions
- Extremes
- Learn robust unknown functions
- Human dynamics or unknown physics
- Correct forcings

Produced by differentiable models



Learning unknown relationships from data (in preparation)



$$R/P_t = (S_m/F_c)^\beta$$




$$R/P_t = ANN(\beta^*, F_c, S_m, S_m/F_c, P_t)$$

Blue line: original power law relation


Red dots: ANN simulations

Black lines: continuous plotting of ANN functions



Example 4. Ecosystem modeling: photosynthesis

Biogeosciences, 20, 2671–2692, 2023
<https://doi.org/10.5194/bg-20-2671-2023>
 © Author(s) 2023. This work is distributed under the Creative Commons Attribution 4.0 License.




(a) Temporal holdout test for the following system

Runs	Corr		RMSE ($\mu\text{mol m}^{-2} \text{s}^{-1}$)		Bias ($\mu\text{mol m}^{-2} \text{s}^{-1}$)		NSE	
	Train	Test	Train	Test	Train	Test	Train	Test
V_{def}+B_{def}	0.565		6.780		1.476		0.041	
V _{def} +B _{def} **	0.592		5.488		1.034		0.318	
V _{def} +B	0.678	0.547	5.887	6.730	1.353	1.754	0.321	-0.084
V+B _{def}	0.769	0.593	4.595	5.677	-0.129	-1.368	0.587	0.229
V+B	0.800	0.748	4.299	4.421	0.037	0.347	0.638	0.532
V+B**	0.774	0.768	4.269	4.198	0.056	0.092	0.597	0.581

** refers to using C3_only plants in dataset


A differentiable, physics-informed ecosystem modeling and learning framework for large-scale inverse problems: demonstration with photosynthesis simulations

Doaa Aboelyazeed¹, Chonggang Xu², Forrest M. Hoffman^{3,4}, Jiangtao Liu¹, Alex W. Jones⁵, Chris Rackauckas⁶, Kathryn Lawcan¹ and Chaoyang Shen¹




Example 4. Ecosystem modeling: photosynthesis



(b) Applied Workflow





Discovering environmental dependencies of previously PFT-dependent parameter



Example 4. Differentiable routing model









MLP n Distribution Trained Against Observed Discharge





Example 6. Fusion of forcings (in preparation)

NLDAS (0.56) > Daymet (0.41) > Maurer (0.03)




Simulation	Forcings	Median NSE	Median KGE	Low flow RMSE (mm/day)	High Flow RMSE (mm/day)
Single forcing w/o bias corr	Daymet	0.737	0.728	0.134	3.990
Multiforcing with bias correction	Daymet, Maurer, NLDAS	0.770	0.780	0.082	3.414

Low bias



Future

- All kinds of models will be differentiable
- Climate change impact assessment will be done using high-quality models that have absorbed big data
- Many theories will be rewritten
- WaterGPT?



Thank you!



@ChaopengShen
cshen@engr.psu.edu

Hydroml.org

<https://mhpi.github.io/benchmarks/>



Shen Multi-scale Hydrology, Processes and Intelligence Group (MHPI)

<http://water.engr.psu.edu/shen/hydroDL.html>

[CUAHSI cyberseminar series on BDML](#)

[WRR special issue on BDML](#)

[AGU Editor's review](#)

Hydrol. Earth Syst. Sci., 22, 5639–5656, 2018
<https://doi.org/10.5194/hess-22-5639-2018>
© Author(s) 2018. This work is distributed under the Creative Commons Attribution 4.0 License.



Hydrology and Earth System Sciences
Open Access
EGU

HESS Opinions: Incubating deep-learning-powered hydrologic science advances as a community

Chaopeng Shen¹, Eric Laloy², Amin Elshorbagy³, Adrian Albert⁴, Jerad Bales⁵, Fi-John Chang⁶, Sangram Ganguly⁷, Kuo-Lin Hsu⁸, Daniel Kifer⁹, Zheng Fang¹⁰, Kuai Fang¹, Dongfeng Li¹⁰, Xiaodong Li¹¹, and Wen-Ping Tsai¹

Water Resources Research

REVIEW ARTICLE

10.1029/2018WR022643

Special Section:

Big Data & Machine Learning in Water Sciences: Recent Progress and Their Use in Advancing Science

A Transdisciplinary Review of Deep Learning Research and Its Relevance for Water Resources Scientists

Chaopeng Shen¹

¹Civil and Environmental Engineering, Pennsylvania State University, University Park, PA, USA

deepLDB

deepLDB -- a mac Landslide database

ARTICLE Check for updates

<https://doi.org/10.1038/s41467-021-26107-z> OPEN

From calibration to parameter learning: Harnessing the scaling effects of big data in geoscientific modeling

Wen-Ping Tsai¹, Dapeng Feng¹, Ming Pan^{2,3}, Hylke Beck⁴, Kathryn Lawson^{1,5}, Yuan Yang^{6,7}, Jiangtao Liu¹ & Chaopeng Shen^{1,5✉}