



Salinity thresholds for understory plants in coastal wetlands

Steven M. Anderson · Emily A. Ury · Paul J. Taillie · Eric A. Ungberg ·
Christopher E. Moorman · Benjamin Poulter · Marcelo Ardón ·
Emily S. Bernhardt · Justin P. Wright

Received: 14 August 2021 / Accepted: 8 November 2021 / Published online: 24 November 2021
© The Author(s), under exclusive licence to Springer Nature B.V. 2021

Abstract The effects of sea level rise and coastal saltwater intrusion on wetland plants can extend well above the high-tide line due to drought, hurricanes, and groundwater intrusion. Research has examined how coastal salt marsh plant communities respond to increased flooding and salinity, but more inland coastal systems have received less attention. The aim of this study was to identify whether ground layer plants exhibit threshold responses to salinity exposure. We used two vegetation surveys throughout the

Albemarle-Pamlico Peninsula (APP) of North Carolina, USA to assess vegetation in a low elevation landscape (≤ 3.8 m) experiencing high rates of sea level rise (3–4 mm/year). We examined the primary drivers of community composition change using Non-metric Multidimensional Scaling (NMDS) and used Threshold Indicator Taxa Analysis (TITAN) to detect thresholds of compositional change based on indicator taxa, in response to potential indicators of exposure to saltwater (Na, and the Σ Ca + Mg) and elevation. Salinity and elevation explained 64% of the variation in community composition, and we found two salinity thresholds for both soil Na⁺ (265 and 3843 g Na⁺/g) and Ca⁺ + Mg⁺ (42 and 126 μ eq/g) where major changes in community composition occur on the APP.

Communicated by Julie C. Zinnert.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s11258-021-01209-2>.

S. M. Anderson · E. A. Ury · E. S. Bernhardt ·
J. P. Wright (✉)
Department of Biology, Duke University, Durham,
NC 27707, USA
e-mail: jw67@duke.edu

S. M. Anderson · C. E. Moorman · M. Ardón
Department of Forestry and Environmental Resources,
North Carolina State University, Raleigh,
NC 27695, USA

P. J. Taillie
Department of Wildlife, Ecology and Conservation,
University of Florida, Gainesville, FL 32611, USA

E. A. Ungberg
Department of Biology, University of North Carolina at
Chapel Hill, Chapel Hill, NC 27599, USA

B. Poulter
Biospheric Science Laboratory, NASA Goddard Space
Flight Center, Greenbelt, MD 20771, USA

Similar sets of species showed sensitivity to these different metrics of salt exposure. Overall, our results showed that ground layer plants can be used as reliable indicators of salinity thresholds in coastal wetlands. These results can be used for monitoring salt exposure of ecosystems and for identifying areas at risk for undergoing future community shifts.

Keywords Coastal wetlands · Saltwater intrusion · Vegetation change · Community composition · Salinity threshold · Regime shifts

Introduction

Coastal wetlands are some of the most vulnerable ecosystems on the frontlines of global change (Craft et al. 2009; Schuerch et al. 2018). As hurricanes and droughts become more common and human infrastructure expands (i.e., canal creation, groundwater extraction for irrigation), the influx of seawater further inland into freshwater-dominated systems (i.e., saltwater intrusion) has become a more severe environmental stressor (Hoepfner et al. 2008; Manda et al. 2014; Zhou et al. 2017; Bhattachan et al. 2018; Herbert et al. 2018). Soil salinization is an increasing threat throughout the world to biologically diverse wetlands that are home to a high number of endemic species which are vulnerable, threatened, or endangered (Herbert et al. 2018; Noss et al. 2015). One of the most dramatic manifestations of salinization is the rapid mortality of trees in freshwater wetlands, leading to the expansion of “ghost forests” (Kirwan and Gedan 2019; Tully et al. 2019; Ury et al. 2021; White et al. 2021). The potential for biodiversity, carbon storage, and plant community change has prompted recent research on the main drivers of changes in freshwater wetlands and brackish marshes (Pezeshki et al. 1990; Pennings et al. 2005; Desantis et al. 2007; Schile et al. 2017; Greene et al. 2019; Taillie et al. 2019a, b; Smart et al. 2020).

Plant species in these coastal landscapes exhibit wide variation in their sensitivity to the magnitude and frequency of salinization (Poulter et al. 2009a, b). Elevated salt exposure causes osmotic and ionic toxicity stress in plants (Kozłowski 1997) and alters their ability to use available nutrients in the soil (Eallonardo et al. 2013). Increased salt exposure

lowers hydraulic conductance, which leads to reducing whole-plant performance (e.g., growth rates), impeding photosynthetic function, causing leaf loss (Chaves et al. 2009; Parida and Das 2005), and limiting regeneration by inhibiting seed germination (Krauss et al. 1998, Kosłowski 1997). Physiological traits govern plants’ ability to tolerate, or adapt to, increased stress (Munns and Tester 2008), and adaptations for tolerating salinity are typically associated with trade-offs against growth or reproduction, so the relative abundance of salt-adapted plants should be associated with the historical regime of salt exposure in an area.

Mechanisms responsible for the sensitivity of individual plant taxa to salinity are well studied (Parida and Das 2005; Parihar et al. 2015); however, comprehensive assessments from field studies that use a quantitative approach for understanding the consequences of species-level sensitivity to salinity on community patterns are sparse in the literature. Often, studies are limited to a small number of species, and the metric of sensitivity is largely based on syntheses or controlled greenhouse experiments that can be limited in scope (Parida and Das 2005; Negrão et al. 2017). In addition, the majority of prior research has focused on forest overstory trees or agricultural crop responses to salinity. Although trees and agricultural crops are ecologically and commercially important, the herbaceous-dominated communities that make up the ground layer of coastal-forested wetlands and the majority of biomass in fresh and saltwater marshes support a high proportion of coastal plain plant diversity (Bratton 1976). In addition to being more diverse, ground layer communities are likely to be more responsive to environmental change than overstory trees because of their shorter life span. Changes in the presence, absence, and relative abundance of ground layer plant taxa may thus serve as an indicator of the abiotic conditions that a community is experiencing and has experienced in the recent past (Osland et al. 2019).

We know plant species respond to salt exposure quite differently; however, one key unknown is whether variability in sensitivity to salt exposure is distributed somewhat continuously across species in an assemblage or if a substantial fraction of species in a regional pool exhibit shifts in abundance at similar points along a gradient. The latter response would suggest that the system can be described as having a

critical threshold or tipping point (Beisner et al. 2003; Scheffer et al. 2012). Identifying these thresholds can be useful for multiple reasons. First, if we can identify critical thresholds, we can detect areas on the landscape that are particularly likely to undergo transitions, and where management interventions might be necessary. Secondly, if monitoring of environmental conditions is difficult or costly, identifying sets of species that are most likely to shift in abundance as thresholds are crossed and focusing monitoring on their presence, absence, or abundance can serve as an alternate tool (i.e., biosensor) for monitoring environmental quality and ecosystem health.

An additional challenge of assessing exposure to marine salts is that while topography and elevation determine vulnerability to flooding and salinity, the vulnerability of ecosystems to salinization is more complex. In addition to rising sea levels along the shoreline (Horton et al. 2009; Kemp et al. 2011), marine salts can penetrate inland well above the high-tide line via storm surge, mixing during drought events, and groundwater intrusion. (Manda et al. 2014; Herbert et al. 2015). Shoreline ecosystems are continuously exposed to marine salts, while in inland wetlands, ecosystem salinization depends upon the retention of salts which can vary due to soil exchange, evaporative concentration, or incorporation into organic matter (Herbert et al. 2015). Furthermore, marine salts are chemically complex and the various constituent ions are likely to be differentially retained and have different effects on vegetation (Tully et al. 2019). Plant responses are integrated responses to salt exposure over time, and previous work on plant responses to saline stress in mature trees found that while changes in tree basal area showed similar declines in response to concentrations of chloride, sulfate, sodium, and elevation, changes in composition were more weakly associated with these drivers (Ury et al. 2020). Given the greater diversity of the plant community in the ground layer and its more rapid turnover time, it seems likely that composition will be affected by the components of salinity, but to what degree, and whether responses are consistent across different components needs further investigation.

The aim of this study was to identify whether there are critical thresholds where large changes in ground layer community composition occur in non-tidal coastal wetlands. Here, we address three questions:

(1) Are potential indicators of exposure to saltwater (elevation, Na, and the Σ Ca + Mg) primary environmental drivers of ground layer plant community composition in coastal wetlands, (2) Are there critical thresholds of salinity exposure that are associated with shifts in composition, and (3) If we identify thresholds, are community responses consistent across multiple components of salinity?

To address these questions, we combined understory, ground layer vegetation data from multiple sources across the Albemarle-Pamlico Peninsula (APP) in eastern North Carolina. Both vegetation and soils data were used to calculate taxa-specific indicator values using a well-known indicator species analytical approach (Dufrene & Legendre 1997) within Threshold Indicator Taxa ANALysis (TITAN,




Fig. 1 Map of the Albemarle-Pamlico Peninsula (approximate centroid coordinates: 35.702451, - 76.245749) with peninsula-wide survey (triangles) and the transect survey (circles) plots colored by Na^+ concentration ($\mu\text{g Na}^+/\text{g}$ soil). Digital elevation model (DEM) of the Albemarle-Pamlico Region is shown in gray color ramp. Water is denoted by light blue areas (Albemarle, Pamlico, and Croatan Sounds surrounding the study area and Atlantic Ocean beyond the Outer Banks), and land outside of the study regions with no DEM in white. See Ury et al. (2020) and Taillie et al. (2019a, b) for additional sampling details

Table 1 Correlation coefficients of each environmental parameter to each axis of the non-metric multidimensional scaling ordination

	Variables	NMDS1	NMDS2	<i>p</i> -value
All sites	Σ Ca⁺ Mg⁺ μeq/g	0.717		*0.01
	Na⁺ μg/g	0.577	0.178	*0.01
	Elevation m	– 0.400	0.176	*0.01
Peninsula-wide survey	Elevation m	0.244	0.211	*0.01
	NH₄⁺ μg/g	0.133	0.170	*0.02
	pH	– 0.125		0.28
	Σ Ca⁺ Mg⁺ μeq/g	– 0.153		0.05
	Mg⁺ μg/g	– 0.153	– 0.122	*0.03
	Na⁺ μg/g	– 0.161	– 0.155	*0.01
	SO₄[–] μg/g	– 0.204		*0.01
	Cl[–] μg/g	– 0.213	– 0.128	*0.01
	NO₃[–] μg/g	– 0.223		*0.02
	Ca ⁺ μ g/g			0.24
	% Soil Moisture			0.54
Transect survey	AFDM			0.57
	Bulk Density g/L			0.74
	Σ Ca⁺ Mg⁺ μeq/g	– 0.148	– 0.497	*0.01
	Na⁺ μg/g	– 0.293	– 0.385	*0.01
	Elevation m			0.33

Bold variables and *p*-values with asterisk are statistically significant (*p* < 0.05)

Results of all three different ordinations (all sites combined, peninsula-wide survey, and transect survey) are presented

Baker and King 2010), a statistical model for detecting thresholds of community change.

Methods

The Albemarle-Pamlico Peninsula (APP) (Fig. 1) is a low-lying region of North Carolina covering ~ 6000 km² with the majority of land area less than 4 m in elevation (Moorhead and Brinson 1995; Poulter et al. 2008) and largely composed of state and federally owned preserves, wildlife refuges, and agricultural lands. Surrounded by the Albemarle, Pamlico, and Croatan Sounds, the APP is the second largest estuarine complex in the USA. The peninsula contains a wide range of woody and herbaceous wetland ecosystems, including upland mesic forests, freshwater-forested wetlands, lowland evergreen and hardwood shrublands, pocosins (poorly drained wetlands with acidic soils, rich in organics), cypress and cedar swamps, and brackish marshes (Brinson 1991). Agricultural fields are hydrologically connected through a network of constructed ditches intended to drain water from higher elevation agricultural field. This ditch network makes the APP particularly vulnerable to salt

stress especially in recent decades (Bhattachan et al. 2018).

Field surveys

To assess the effects of salinity on herbaceous wetland vegetation in the APP, we used two data sets. The first data set (the “transect” data set) was originally established in 2004 to investigate vegetation change across a salinity gradient (Poulter 2005), but was then resampled during 2016/2017 to investigate changes related to rising sea level (Taillie et al. 2019a, b). The original study used stratified random sampling to measure vegetation at five public properties throughout the peninsula (Fig. 1). In addition to the stratification by geography (i.e., across properties), the sampling was stratified by vegetation community, where each property was delineated into three zones: intact forest, transition/ghost forest, and brackish marsh, according to aerial photography and visual inspection in the field. Within each vegetation community at each site, seven 11-m-radius plots were randomly established. However, the marsh vegetation community was absent from one property, which resulted in 98 total plots. Rather than inventory understory vegetation across the entire 11-m-radius

Table 2 Environmental thresholds (i.e., change points) for three salinity gradients and range of observed values

	Elevation m (− 0.1 to 3.76)	Na ⁺ µg/g (75 to 9329)	∑ Ca ⁺ and Mg ⁺ µeq/g (0.64 to 288.9)
Decreasing (z−) taxa Threshold (# of plots)	0.24 (169)	265 (124)	42.01 (186)
Increasing (z+) taxa Threshold (# of plots)	1.6 (32)	3843 (465)	261.3 (285)
Prop. of pure and reliable taxa	0.34	0.66	0.62
No. non-responsive taxa	56	41	43

The number of plots below (z−) or above (z+) the community change points for each environmental gradient is in parentheses. We observed a total of 112 taxa, 30 of which were rare (< 3 observations). The proportion of pure and reliable taxa and number of non-responding (not pure and reliable) taxa are included

plot, 5 and 1-m² sub-plots were established, within which understory vegetation cover was estimated. The percent cover for each species was then averaged across the 5 sub-plots to obtain an estimated percent cover for each species identified within each plot. Only the data collected during the 2016 survey were used in this study. See Poulter (2005) and Taillie et al. (2019a, b) for additional information regarding the transect survey protocol.

The second data set (the “peninsula-wide” data set) has not previously been published and focused on sampling a broader gradient of wetland types and environmental conditions in 2016. In total, vegetation surveys were conducted at 34 sites (5 sites from the transect survey and 29 sites from the peninsula-wide survey) in 669, 1-m² plots to assess the range of environmental conditions where species occur throughout the APP (Fig. 1). The peninsula-wide data set followed the Carolina Vegetation Survey protocol (CVS, Peet et al. 1998). We delineated 29 sites that were 400 m² or 100 m² depending on the heterogeneity of dominant vegetation. At each site, we systematically located two (in the 100-m² sites) or eight (in the 400-m² sites) 1-m² plots for sampling herbaceous vegetation (woody and non-woody < 2 m tall). We estimated percent cover of all species located in these plots. In both data sets, we identified all plants rooted in each plot less than 1 m in height and estimated areal cover abundances for each species.

Soils from the transect survey were taken from the top 10 cm, sieved (2-mm mesh), and analyzed major cation using the Mehlich III extraction (Brookside Labs., New Knoxville, Ohio). At each site in the

peninsula-wide survey, we sampled soils to 10-cm depths and sieved through 2-mm mesh; extracted fresh soils with deionized water; and analyzed for ion concentrations including base cations (Ca²⁺, Mg²⁺, and Na⁺) on a CS12A column, and Cl[−] using a AS-18 analytical column with Dionex ICS 2000 (Dionex Corporation, Sunnyvale, CA). Because two different soil extraction methods were used in the two different surveys (peninsula-wide and transect surveys), we included methodology as categorical factor within a second set of ordinations to test whether extraction method had a significant contribution to the outcome of the environmental drivers. We used a LiDAR-derived digital surface model (DSM) raster data set with 6.1-m resolution and vertical accuracy of ± 0.13 m for calculating the elevation at each site analyzed in the study. This DSM was part of a statewide data set collected in the 1990s. (See Bhattachan et al. 2018 for additional information.) We calculated the average elevation at each site since microtopography varied across 4–18 points that were at least 10 m apart at each location.

Plant community responses

To determine whether salinity is a major driver of species composition throughout the APP, we used non-metric multidimensional scaling (NMDS), a commonly used ordination method for plant community assessment. We used the R version 3.6.3 (R Core Team 2019) packages *ecodist* (Goslee and Urban 2007) and *vegan* (Oksanen et al. 2011) to run the ordination analyses. We focused on three potential




Fig. 2 Frequency distributions and correlations of all three environmental parameters (elevation, Na^+ , and the ΣCa^+ and Mg^+) hypothesized to be the main drivers of understory community composition and used later in the threshold indicator

indicators of salinization: elevation, Na^+ concentrations, and the sum of Ca^+ and Mg^+ . Elevation is a variable that is readily available in all contexts and is frequently thought of as a proxy for exposure due to flooding and sea level rise, although it may not perfectly predict the risk (Bhattachan et al. 2018) and impacts (Ury et al. 2020) of exposure. Sodium and chloride are the most abundant ions in seawater and we focused Na^+ as a metric of salinization because we did

not have Cl^- concentrations from the transect survey data set. To confirm that Na^+ would be a good indicator of saltwater intrusion, we tested the correlation of Na^+ and Cl^- (the most abundant ions in seawater) in soils from the peninsula-wide survey of the APP (Figure S1, $r^2 = 0.976$). Finally, in addition to changes in the ionic strength of soils, one of the major impacts of salinization is through altering the alkalinity of soils (Tully et al. 2019). To assess this impact,

not have Cl^- concentrations from the transect survey data set. To confirm that Na^+ would be a good indicator of saltwater intrusion, we tested the correlation of Na^+ and Cl^- (the most abundant ions in seawater) in soils from the peninsula-wide survey of the APP (Figure S1, $r^2 = 0.976$). Finally, in addition to changes in the ionic strength of soils, one of the major impacts of salinization is through altering the alkalinity of soils (Tully et al. 2019). To assess this impact,

Table 3 Change point values for pure and reliable taxa identified by TITAN as salt sensitive/decreasing taxa (–) or tolerant/increasing taxa (+) for Na⁺ µg/g and ∑ Ca⁺ & Mg⁺ µeq/g gradients and sensitive/increasing taxa (+) or tolerant/decreasing taxa (–) along the elevation gradient

Taxa	Observations	Elevation m	Na ⁺ µg/g	∑ Ca ⁺ + Mg ⁺ µeq/g
<i>Acer rubrum</i> (ACRU)	105	3.143 ⁺		6.13 ⁻
<i>Alternanthera philoxeroides</i> (ALPH)	7		293 ⁻	20.30 ⁻
<i>Andropogon</i> spp. (ANSP)	7		566 ⁻	81.79 ⁻
<i>Arundinaria gigantea</i> (ARGI)	9	0.196 ⁻	293 ⁻	76.20 ⁻
<i>Arundinaria tecta</i> (ARTE)	12	0.724 ⁺	101 ⁻	2.06 ⁻
<i>Asimina triloba</i> (ASTR)	6	3.757 ⁺	92 ⁻	3.63 ⁻
<i>Baccharis halimifolia</i> (BAHA)	49			6.13 ⁺
<i>Bignonia capreolata</i> (BICA)	5		265 ⁻	3.63 ⁻
<i>Boehmeria cylindrica</i> (BOCY)	4	3.757 ⁺	92 ⁻	
<i>Callicarpa americana</i> (CAAM)	4	0.610 ⁺	95 ⁻	1.04 ⁻
<i>Campsis radicans</i> (CARA)	7	3.207 ⁺	100 ⁻	73.89 ⁻
<i>Carex</i> spp. (CASP)	9		263 ⁻	3.63 ⁻
<i>Carpinus caroliniana</i> (CACA)	7		502 ⁻	1.04 ⁻
<i>Chasmanthium laxum</i> (CHLA)	6			12.10 ⁻
<i>Cladium jamacense</i> (CLJA)	218	0.137 ⁻	626 ⁺	111.93 ⁺
<i>Clethra alnifolia</i> (CLAL)	5		76 ⁻	0.98 ⁻
<i>Distichlis spicata</i> (DISP)	73	0.307 ⁻	3,581 ⁺	227.81 ⁺
<i>Gelsemium sempervirens</i> (GESE)	12		347 ⁻	73.81 ⁻
<i>Hydrocotyle verticillata</i> (HYVE)	23		1,165 ⁻	
<i>Ilex coriacea</i> (ILCO)	8		796 ⁻	1.35 ⁻
<i>Ilex glabra</i> (ILGL)	43	0.565 ⁺	681 ⁻	104.34 ⁻
<i>Ilex opaca</i> (ILOP)	9	1.376 ⁺	211 ⁻	3.63 ⁻
<i>Iva frutescens</i> (IVFR)	13		1,165 ⁺	
<i>Juncus roemerianus</i> (JURO)	67		1,165 ⁺	98.58 ⁺
<i>Juncus</i> sp. (JUSP)	9	0.610 ⁺	953 ⁻	91.09 ⁻
<i>Kosteletzkya pentacaros</i> (KOPE)	21	0.518 ⁻	543 ⁻	81.55 ⁺
<i>Lemma</i> sp. (LESP)	8		307 ⁻	4.59 ⁻
<i>Liquidambar styraciflua</i> (LIST)	6		508 ⁻	
<i>Liriodendron tulipifera</i> (LITU)	11	1.612 ⁺	211 ⁻	3.81 ⁻
<i>Lyonia lucida</i> (LYLU)	54	0.305 ⁺	690 ⁻	104.34 ⁻
<i>Microstegium vimineum</i> (MIVI)	3	3.482 ⁺	94 ⁻	
<i>Mitchella repens</i> (MIRE)	17		269 ⁻	3.56 ⁻
<i>Morella cerifera</i> (MOCE)	120		1253 ⁻	
<i>Nyssa biflora</i> (NYBI)	6		282 ⁻	
<i>Osmunda regalis</i> (OSRE)	78			51.10 ⁺
<i>Parthenocissus quinquefolia</i> (PAQU)	33		269 ⁻	71.51 ⁻
<i>Persea palustris</i> (PEPA)	93		769 ⁻	85.58 ⁻
<i>Persicaria</i> spp. (PESP)	3		1,007 ⁻	46.33 ⁻
<i>Phragmites australis</i> (PHAU)	81	0.120 ⁻	656 ⁺	61.35 ⁺
<i>Pinus serotina</i> (PISE)	17		653 ⁻	99.65 ⁻
<i>Pinus taeda</i> (PITA)	44	0.437 ⁺	269 ⁻	0.98 ⁻
<i>Prunus</i> sp. (PRSP)	3			1.04 ⁻
<i>Quercus nigra</i> (QUNI)	13		267 ⁻	3.63 ⁻
<i>Rubus</i> spp. (RUSP)	15		617 ⁻	77.51 ⁺
<i>Sagittaria latifolia</i> (SALA)	9	0.305 ⁻	138 ⁻	70.56 ⁻
<i>Saururus cernuus</i> (SACE)	8		100 ⁻	3.15 ⁻
<i>Schoenoplectus americana</i> (SCAM)	50	0.309 ⁻	4,747 ⁺	243.86 ⁺
<i>Smilax</i> sp. (SMSP)	116	1.260 ⁺	873 ⁻	81.30 ⁻
<i>Spartina alterniflora</i> (SPAL)	11	0.021 ⁻	3,749 ⁺	262.37 ⁺
<i>Spartina patens</i> (SPPA)	57	0.476 ⁻	347 ⁺	
<i>Sphagnum</i> sp. (SPHSP)	7	0.380 ⁺	94 ⁻	44.27 ⁻
<i>Symplocos tinctoria</i> (SYTI)	7		98 ⁻	1.44 ⁻
<i>Taxodium distichum</i> (TADI)	5	3.080 ⁺	98 ⁻	3.15 ⁻
<i>Toxicodendron radicans</i> (TORA)	229		2,470 ⁻	6.13 ⁺
<i>Typha latifolia</i> (TYLA)	12			42.01 ⁻
<i>Vaccinium</i> spp. (VASP)	9		681 ⁻	88.29 ⁻
<i>Vitis rotundifolia</i> (VIRO)	25	0.456 ⁺	169 ⁻	45.34 ⁻
<i>Woodwardia aereolata</i> (WOAE)	16		263 ⁻	3.63 ⁻
<i>Woodwardia virginica</i> (WOVI)	116		1,175 ⁻	118.82 ⁻

Table 3 continued

Shading corresponds the classification TITAN assigned to each species as tolerant (dark gray) or sensitive (light gray) along the particular salinity gradient

we looked at the sum of Ca^+ and Mg^+ . We used vector fitting (vf) in *ecodist* (Goslee and Urban 2007) to display all environmental variables included in the ordination analysis to further describe the main drivers of species composition throughout the study area (Oksanen et al. 2011; Ury et al. 2020). Prior to analyses, species abundances were relativized and all environmental variables were tested for normality using the Shapiro–Wilk Test. Because of potential differences in methodology and ranges of environmental variables across the two data sets, we conducted three separate NMDS ordinations on cover data: (1) using the combined data sets from both surveys; (2) using only the data from the “transect” data set; and (3) using only the data from the “peninsula wide” data set.

Species responses

To assess whether there were community-level thresholds in ground level species responses to the salinization gradients, we used the TITAN (version 2.1, Baker et al. 2015, King et al. 2011) package in R (R Core Team 2019). TITAN identifies change points in abundance (i.e., thresholds) for each individual taxa along each environmental gradient tested. We categorized taxa as negative/decreasing (z^-) or positive/increasing (z^+) responders to the three environmental gradients associated with salinization. We used a plot by species matrix for all taxa that occurred at ≥ 3 plots throughout the study area to ensure that species change points could be estimated.

Based on species abundance and frequency in the data, each species was given a taxa-specific indicator value (IndVal) for the indicator species analysis (Dufrêne & Legendre 1997). By relativizing the species abundance data, TITAN assigned z scores to each species z^- or z^+ responders, which were summed for community-level analyses. At the community level, TITAN can identify two thresholds along the gradient each corresponding to negative or positively responding taxa. Bootstrap replication ($n = 500$) was

used to estimate uncertainty in the change points. We only reported results from pure taxa (change in abundance and frequency of a given species was in the same positive or negative direction along the gradient for $> 95\%$ of the bootstrap replicate runs) and reliable taxa [$> 95\%$ of bootstrap replicate runs needed to be significantly different from a random distribution of ($p \leq 0.05$)]. We used filtered z -score values which only include pure and reliable taxa in the model.

Results

Environmental variables associated with soil salinity

Mean elevation of plots was 0.51 m (SD 0.65) with 91.6% of plots below 1 m (Fig. 2), but the two data sets had different means (“transect” = 0.32 ± 0.20 m; “peninsula wide” = 0.96 ± 1.02 m). The distribution of soil Na^+ concentrations was right skewed with a mean concentration of 1718 $\mu\text{g Na}^+/\text{g soil}$ (SD 1891.3, Fig. 2). The sum of water extractable $\text{Ca}^+ + \text{Mg}^+$ was bimodal with a peak in plots below 25 $\mu\text{eq Ca}^+\text{Mg}^+/\text{g soil}$ and a mean concentration of 106.7 $\mu\text{eq Ca}^+\text{Mg}^+/\text{g soil}$ (SD 82.6, Fig. 2). Mean $\text{Ca}^+ + \text{Mg}^+$ concentrations differed across the two surveys (transect survey = 145.1 ± 65.44 m; peninsula-wide survey = 13.86 ± 27.56 m). Sodium and $\text{Ca}^+ + \text{Mg}^+$ were significantly correlated ($p = < 0.005$, $r^2 = 0.497$, Fig. 2). We observed higher ionic strength values (mean = 1927.4 $\text{g Na}^+/\text{g}$) in plots less than 1 m in elevation and lower salinity (mean = 106.7 $\text{g Na}^+/\text{g}$) in plots greater than 1 m in elevation (Fig. 2). Low elevation plots also exhibited large variance in salinity values.

Effects of salinization on community composition

We identified 112 plant species in the study area and 84 species occurred in more than three plots across the study. Of those species identified in both surveys, 25 species occurred in at least 5% of the plots and were included in the NMDS ordinations (stress = 0.179). Sodium, $\text{Ca}^+ + \text{Mg}^+$, and elevation collectively explained 64% of the variability in community composition in the ordination combining both surveys (Fig. 3). NMDS1 explained 32.3% of the variability and all three salinity variables (Na^+ , $\text{Ca}^+ + \text{Mg}^+$, and elevation) were significant drivers of this axis ($p = 0.01$) (Table 1). NMDS2 explained 31.7% of the variation and was driven by Na^+ and elevation ($r = 0.178$ and 0.176) (Table 1). Because the distributions of these variables differed between the two

surveys, we also ran ordinations for each survey separately. For the transect survey, Na^+ and $\text{Ca}^+ + \text{Mg}^+$ were significant drivers on both NMDS1 and NMDS2 ($r = -0.15$ – 0.50 , $p = 0.01$), but not elevation. For the peninsula-wide survey, elevation was a significant predictor of community composition ($p = 0.01$) as was Cl^- , Ca^+ , and Mg^+ ($p = 0.01$ – 0.03) (Table 1). Together, these analyses show that soil Na^+ was the strongest driver of community composition across all plots in both surveys. $\text{Ca}^+ + \text{Mg}^+$ was also a significant driver across all plots, but only marginally significant in the peninsula-wide survey, although Mg^+ was also significant (Table 1, Figure S3). On the other hand, elevation was a significant driver of composition in the peninsula-wide survey, but was not significant when only considering the lower elevations (< 1 m above




Fig. 3 Non-metric multidimensional scaling (NMDS) ordination analysis of all sites with three significant environmental variables fit as vectors ($p < 0.05$). Sodium, elevation, calcium, and magnesium explains 64% herbaceous plant community composition in two axes. Each plot is indicated by solid gray

circles. Taxa names are weighted average abundance based on species presence. The length of the arrow indicates a large degree of statistical significance. See Table 1 for statistics of NMDS1 and NMDS2, and see Table 3 for taxa full names

sea level) as seen in the transect survey (Table 1, Figure S2).

Threshold indicator taxa analysis (TITAN)

TITAN retained 54 pure and reliable taxa ($p \leq 0.05$, 66% of all taxa) along the Na^+ gradient. Forty-three of those taxa were categorized as decreasing (z^-) species and 11 as increasing (z^+) species accounting for 81.1% and 18.9% of the taxa along the Na^+ gradient, respectively (Table 2, Fig. 4). Thresholds for individual decreasing species were between 75 and 3017 $\mu\text{g Na}^+/\text{g}$ and the summed z^- community threshold (the point along the gradient at which the largest number of sensitive taxa were declining in abundance) was

265 $\mu\text{g Na}^+/\text{g}$. Thresholds for individual increasing taxa (z^+) were between 353 and 4388 $\mu\text{g Na}^+/\text{g}$ and the summed community threshold was 3843 $\mu\text{g Na}^+/\text{g}$ (Table 2, Fig. 4).

Results using marine base cations (Ca^+ and Mg^+) were similar to the Na^+ gradient analysis. TITAN retained 51 reliable taxa (or 62% of all taxa) along this gradient. Forty decreasing (z^-) and 11 increasing (z^+) species accounted for 78.4% and 21.6% of taxa along the $\text{Ca}^+ + \text{Mg}^+$ gradient, respectively (Table 3, Figure S4). Thresholds for individual decreasing taxa (z^-) along the $\text{Ca}^+ + \text{Mg}^+$ gradient occurred between 0.98 and 110.6 $\mu\text{eq/g}$ with a community threshold at 42.01 $\mu\text{eq/g}$. Increasing taxa (z^+) thresholds occurred between 62.9 and 264.8 $\mu\text{eq/g}$ with a




Fig. 4 Sharp decline in the majority of taxa below 2000 $\mu\text{g/g}$. Negative (gray) and positive (black) responding taxa (on y-axis) from TITAN along the Na^+ gradient. Gray vertical line at 265 $\mu\text{g/g}$ denotes the summed z^- -scores for negative responding (z^-) community threshold, and the dark gray vertical line at 3843 $\mu\text{g/g}$ denotes the summed z^+ -scores for positive responding (z^+) community threshold. Letters next to each taxa

abbreviation relate to whether it is a tree, shrub, forb, or graminoid species as noted in the legend. Points indicate the change point (or threshold) for each species sized by its indicator value (IndVal) which corresponds to individual species contribution to the summed z^- -score/community-level threshold. Horizontal lines for each species indicate the 5–95% quartile range of that taxa

summed community threshold at 126.31 $\mu\text{g/g}$ (Table 2, Figure S5).

Fewer taxa were retained for the TITAN analysis of responses to the elevation gradient (30 species); 17 taxa were categorized as decreasing/tolerant (z^-) and 10 as increasing/sensitive (z^+), accounting for 66.7% and 33.3% of total retained taxa, respectively (Table 3, Figure S5). In this TITAN analysis, increasing taxa (higher elevation, less salt tolerant) occurred between 0.29 and 3.76 m with the summed community threshold (z^+) at 1.61 m and decreasing taxa (lower elevation/more salt tolerant) between 0.01 and 0.53 m with a summed community threshold (z^-) at 0.24 m (Table 2, Figure S5).

Of the three gradients we analyzed in TITAN, Na^+ and $\text{Ca}^+ + \text{Mg}^+$ had the most taxa in common (65.4%) and elevation had 51.9% of species in common with Na^+ and 48.1% with $\text{Ca}^+ + \text{Mg}^+$ (Table 3). However, in some cases TITAN classified decreasing and increasing species differently among all three environmental gradients (Table 3). For example, *Sagittaria latifolia* was classified as a decreasing species as Na^+ or $\text{Ca}^+ + \text{Mg}^+$ increased, but as an increasing species as elevation increased. Similarly, *Toxidendron radicans* was classified as a decreasing species on the Na^+ gradient, but an increasing species on the $\text{Ca}^+ + \text{Mg}^+$ gradient (Table 3).

Discussion

This study sought to identify whether there are critical thresholds where changes in ground layer community composition occur in non-tidal coastal wetlands in response to salinity exposure. Here, we combined NMDS and TITAN analyses to reveal a strong association between soil salinity and elevation and plant community composition, with the majority of ground layer species (herbaceous and woody) responding negatively to higher levels of salinity. Therefore, our analyses confirm that salinity is a primary environmental driver of understory plant community composition in coastal wetlands and suggest that there is a narrow range of salt concentrations where most sensitive taxa are lost. Specifically, when soil salinity exceeded 265 $\mu\text{g Na}^+/\text{g}$, 44% of species were lost. Our results also indicated that species responses were largely congruent across

multiple indicators of salinity exposure (Na^+ and $\text{Ca}^+ + \text{Mg}^+$) and elevation, although responses to elevation were not as consistent. We revealed a number of reliable indicator taxa that may be useful as early warning signals of salinization in freshwater wetlands in the coastal plain of the southeastern USA.

Associations of community composition and environmental drivers

As predicted, the ground layer plant communities in this landscape were structured by factors associated with salt exposure. We looked at metrics of risk of exposure (elevation), the effects of ionic strength (Na^+), and the effects of changes to base cations ($\text{Ca}^+ + \text{Mg}^+$) and determined all three vectors were correlated with community composition. However, in this remarkably flat landscape, differences in elevation proved to be a less important driver of community composition than measured salinity impacts, consistent with Taillie et al. (2019a, b, i.e., the transect survey). It was only when including the relatively higher elevation plots present in the peninsula-wide survey that we were able to detect an impact of elevation on composition. Salinity (Na^+ and $\text{Ca}^+ + \text{Mg}^+$) concentrations were extremely variable in plots that were < 1 m elevation and not as variable in higher elevation plots > 1 m elevation. Nevertheless, salinity-associated variables were significant predictors of composition across the entire data set as well as in each of the individual data sets. Local microtopography and slope are strong drivers of plant occurrence when studying low-lying coastal wetland ecosystems (Moleno et al. 2021). Elevation can be an important driver of species presence or absence, but it becomes more important with increasing geographic scale, while hydrology and salinity typically appear to be more influential as local-scale drivers on plant communities in coastal ecosystems (Herbert et al. 2015).

Reliable indicators of salinity thresholds

This study applied formalized indicator taxa analysis (e.g., TITAN) for assessing sensitivity of coastal wetland plants to salinity stress. We were able to identify community thresholds based on where the greatest change in abundance and frequency of understory plant taxa occurred along the soil salinity gradient. As we expected, taxa most sensitive to soil

salinity were those typical of higher elevation forest types, dominated by seedlings and saplings of deciduous hardwoods (e.g., *Acer rubrum*, *Carya ovata*, *Clethra alnifolia*, *Asimina triloba*, and *Liriodendron tulipifera*) and evergreens trees and shrubs (*Pinus taeda* and *Ilex opaca*). These species typically have limited exposure to seawater except for large hurricanes which can deliver seawater further inland to higher elevation sites (Middleton 2016). However, adults of species like *P. taeda* and *P. serotina* are dominant in coastal freshwater wetlands and tolerate periodic pulses of salts at low concentrations (Poulter et al. 2008) and as seedlings have been shown to increase aboveground biomass in response to salt exposure (Anderson 2020). We also identified salt-sensitive herbaceous indicator taxa characteristic of upland forests (e.g., *Arundinaria tecta*, *Clethra alternifolia*, *Mitchella repens*) and salt-tolerant taxa common to brackish conditions: grasses (e.g., *Distichlis spicata*, *Sporobolus* (formerly *Spartina*) *alterniflora*, *Spartina patens*) and sedges (e.g., *Cladium jamaicense*, *Scirpus americanus*).

We identified a particular range of salinity concentrations between the summed z^- and z^+ community threshold ($\text{Na}^+ = 265\text{--}3843 \mu\text{g/g}$, $\text{Ca}^+ + \text{Mg}^+ = 42.01\text{--}126.31 \mu\text{eq/g}$, elevation = 0.24–1.61 m) where taxa are likely to be good indicators, or early warning signs, of projected vegetation change. It is in this range where the abundance of decreasing (z^-) and increasing (z^+) taxa along the salinity gradient overlap, reminiscent of pre-existing or currently expanding transitional/“ghost” forests in the transect survey (Fig. 5). Sites in this range of salinity values had a mix of salt sensitive and salt-tolerant herbaceous layer taxa, some with live mature trees and others with dead trees. For example, we documented transitioning forests dominated by evergreen saplings (e.g., *Persea palustris* and *Ilex glabra*) and rhizomatous shrubs (e.g., *Lyonia lucida*). These species have relatively high salt sensitivities and form dense shrub-scrub thickets where relatively salt-tolerant fern species (e.g., *Osmunda regalis*, *Woodwardia virginica*) are common. We found that other species that occur within highly saline regions along the salinity gradient beyond the z^- threshold include invasive generalists (e.g., *Phragmites australis*), natives often associated with disturbed, or recovering, forests from salt, inundation, or fire exposure (e.g., *Morella cerifera*, *Baccharis halimifolia*, *Iva frutescens*, *Rubus* spp., and




Fig. 5 Conceptual model of changes in plant communities controlled by soil salinization and elevation/flooding from sea level rise and climate change as seen in NMDS analyses. Examples of indicator species, decreasing (z^-) and increasing (z^+) taxa from the TITAN analyses characteristic of forest (yellow), transition (pink), or marsh (purple) communities. This model builds on the model presented by Fagherazzi et al. (2019)

T. radicans), as well as salt-tolerant freshwater marsh species (e.g., *Juncus roemarianus*, *Kosteletzkya pentacarpos*). It is important to note that our results reflect a snapshot approach despite the potentially dynamic nature of soil salinization over space and time throughout the coastal plain (Tully et al. 2019). The overall landscape dynamics suggest that these associations may be unstable (Smart et al. 2020), and we predict that areas once dominated by forest and freshwater marsh plant species known to have a competitive advantage (Crain et al. 2004) become intolerant to salty conditions and converted to shrub-scrub and ghost forests. This pattern has been documented in Palmetto-Peartree Preserve on the Albemarle-Pamlico Peninsula between 1996 and 2017 (White et al. 2021) and will eventually be colonized by salt-tolerant marsh species (Crain et al. 2004; Sutter et al. 2015).

All three gradients (Na^+ , $\text{Ca}^+ + \text{Mg}^+$, and elevation) were drivers of community composition and they each identified two critical salinity thresholds for each species; however, species salinity thresholds were not all consistent across the gradients. We documented consistent responses in 19 species across all three

gradients, consistent responses of 22 species between Na^+ and elevation, and consistent responses of 21 species between elevation and $\text{Ca}^+ + \text{Mg}^+$. Most strikingly, 42 species had consistent responses between Na^+ and $\text{Ca}^+ + \text{Mg}^+$, which suggest that elevation alone is not as good a predictor of exposure to salinity because marine salts can be delivered, concentrated, or leached by hydrologic droughts, hurricanes, and storm surges (Tully et al. 2019). It is likely that a large proportion of coastal wetland communities experience salinities in this range over short time periods, but long-term exposure is more episodic.

This study presents a compelling case for characterizing salinity thresholds by conducting indicator taxa analyses of understory plant communities. We demonstrated that Na^+ , $\text{Ca}^+ + \text{Mg}^+$, and elevation are main abiotic drivers of community composition. We were able to define salinity thresholds for the majority of taxa identified in the region and subsequently two community-level thresholds for each salinity gradient. The study demonstrates the reliability of combining ordination and threshold indicator taxa analyses as a quantitative approach to understanding the vulnerability of plant communities to salinity and other stressors. Moreover, TITAN is a promising tool for identifying understory plant communities that are likely to experience shifts as a result of increasing saltwater intrusion events. Our work presents an additional use of TITAN for plant community assessment across a naturally occurring gradient. These techniques should continue to be tested for multiple stressors across other habitat types and be used more widely for management and conservation efforts of vulnerable ecosystems over space and time.

Acknowledgements The authors would like to sincerely thank the hard work and dedication of Ethan Baruch, Christina Bergemann, Charles Camp, Ashley Green, Sam Mahanes, Matthew Stillwagon, and Anna Wade in data collection in the field and lab. They would also like to thank the members of the Bernhardt lab at Duke University, the Ardon lab at N.C. State University, and Justin T. Mann at Binghamton University for crucial feedback on various editions of this manuscript, Robert Peet and the Carolina Vegetation Survey for access to survey plots, and Bonnie McGill, PhD for plant illustrations in Fig. 4. Sincere thanks to National Science Foundation for funding this research and earlier sources of funding (from a NASA Earth Science Fellowship) that established the plot networks on the Albemarle Peninsula. Additional thanks to the College of Natural Resources at North Carolina State University and Department of the Interior

Southeast Climate Adaptation Science Center and for funding data collection efforts for the transect survey.

Author contributions JW, EB, CM, and MA procured funding to support this research; SA, JW, EB, and EU conceived the primary research idea; SA, PT, and EU collected field data; SA, BP, and PT selected research sites; SA conducted all analysis and wrote the manuscript with JW; EU generated maps; and all authors discussed results and contributed in edits to manuscript.

Funding National Science Foundation (Coastal SEES Collaborative Research Award Grant No. 1426802, DEB-1713435, and DEB 1713502).

Data availability The data from the peninsula-wide survey are archived and openly available for download via Dryad (<https://datadryad.org/>) and data from the transect survey are available via Taillie and Poulter (2019) via Pangaea (<https://doi.pangaea.de/10.1594/PANGAEA.896941>).

Code availability Code is available on request from the lead author.

Declarations

Conflict of interest The authors declare that they have no conflict of interest.

Consent to participate Not applicable.

Consent for publication Not applicable.

Ethical approval Not applicable.

Nomenclature A.S. Weakley 2015, Flora of the Southern and Mid-Atlantic States.

References

- Anderson, S. M. (2020). Physiological responses of coastal freshwater wetland trees to saltwater intrusion: a greenhouse mesocosm experiment. M.Sc. Thesis, North Carolina State University. Raleigh, NC
- Baker ME, King RS (2010) A new method for detecting and interpreting biodiversity and ecological community thresholds. *Methods Ecol Evol* 1(1):25–37. <https://doi.org/10.1111/j.2041-210x.2009.00007.x>
- Baker ME, King RS, Kahle D (2015) TITAN2: Threshold indicator taxa analysis. R package version 2.1. Retrieved from <https://CRAN.R-project.org/package=TITAN2>
- Beisner BE, Haydon DT, Cuddington K (2003) Alternative stable states in ecology. *Front Ecol Environ* 1(7):376–382. [https://doi.org/10.1890/1540-9295\(2003\)001\[0376:ASSIE\]2.0.CO;2](https://doi.org/10.1890/1540-9295(2003)001[0376:ASSIE]2.0.CO;2)
- Bhattachan A, Emanuel RE, Ardon M, Bernhardt ES, Anderson SM, Stillwagon MG, Ury EA, BenDor TK, Wright JP (2018) Evaluating the effects of land-use change and future

- climate change on vulnerability of coastal landscapes to saltwater intrusion. *Elem Sci Anth* 6(1):62. <https://doi.org/10.1525/elementa.316>
- Bratton SP (1976) Resource division in an understory herb community: responses to temporal and microtopographic gradients. *Am Nat* 110(974):679–693. <https://doi.org/10.1086/283097>
- Brinson MM (1991) Landscape properties of pocosins and associated wetlands. *Wetlands* 11(1):441–465. <https://doi.org/10.1007/BF03160761>
- Chaves MM, Flexas J, Pinheiro C (2009) Photosynthesis under drought and salt stress: regulation mechanisms from whole plant to cell. *Ann Bot* 103:551–560. <https://doi.org/10.1093/aob/mcn125>
- Craft C, Clough J, Ehman J, Joye S, Park R, Pennings S, Guo H, Machmuller M (2009) Forecasting the effects of accelerated sea-level rise on tidal marsh ecosystem services. *Front Ecol Environ* 7(2):73–78. <https://doi.org/10.1890/070219>
- Crain CM, Silliman BR, Bertness SL, Bertness MD (2004) Physical and biotic drivers of plant distribution across estuarine salinity gradients. *Ecology* 85(9):2539–2549. <https://doi.org/10.1890/03-0745>
- Desantis LRG, Bhotika S, Williams K, Putz FE (2007) Sea-level rise and drought interactions accelerate forest decline on the Gulf Coast of Florida, USA. *Glob Change Biol* 13(11):2349–2360. <https://doi.org/10.1111/j.1365-2486.2007.01440.x>
- Dufrène M, Legendre P (1997) Species assemblages and indicator species: the need for a flexible asymmetrical approach. *Ecol Monogr* 67(3):345–366. <https://doi.org/10.2307/2963459>
- Eallonardo AS, Leopold DJ, Fridley JD, Stella JC (2013) Salinity tolerance and the decoupling of resource axis plant traits. *J Veg Sci* 24(2):365–374. <https://doi.org/10.1111/j.1654-1103.2012.01470.x>
- Fagherazzi S, Anisfeld SC, Blum LK, Long EV, Feagin RA, Fernandes A, Kearney WS, Williams K (2019) Sea level rise and the dynamics of the marsh-upland boundary. *Front Environ Sci*. <https://doi.org/10.3389/fenvs.2019.00025>
- Goslee SC, Urban DL (2007) The ecodist package for dissimilarity-based analysis of ecological data. *J Stat Softw* 22(7):1–19. <https://doi.org/10.18637/jss.v022.i07>
- Greene RE, Evans KO, Gray MT, Jones-Farrand DT, Wathen WG (2019) Using a coproduction approach to map future forest retention likelihood in the southeastern United States. *J for*. <https://doi.org/10.1093/jofore/fvz063>
- Herbert ER, Boon P, Burgin AJ, Neubauer SC, Franklin RB, Ardón M, Hopfensperger KN, Lamers LPM, Gell P (2015) A global perspective on wetland salinization: ecological consequences of a growing threat to freshwater wetlands. *Ecosphere* 6(10):206. <https://doi.org/10.1890/ES14-00534.1>
- Herbert ER, Schubauer-Berigan J, Craft CB (2018) Differential effects of chronic and acute simulated seawater intrusion on tidal freshwater marsh carbon cycling. *Biogeochemistry* 138(2):137–154. <https://doi.org/10.1007/s10533-018-0436-z>
- Hoepfner SS, Shaffer GP, Perkins TE (2008) Through droughts and hurricanes: tree mortality, forest structure, and biomass production in a coastal swamp targeted for restoration in the Mississippi River Deltaic Plain. *For Ecol Manage* 256(5):937–948. <https://doi.org/10.1016/j.foreco.2008.05.040>
- Horton BP, Peltier WR, Culver SJ, Drummond R, Engelhart SE, Kemp AC, Mallinson D, Thieler ER, Riggs SR, Ames DV, Thomson KH (2009) Holocene sea-level changes along the North Carolina coastline and their implications for glacial isostatic adjustment models. *Quat Sci Rev* 28(17–18):1725–1736
- Kemp AC, Horton BP, Donnelly JP, Mann ME, Vermeer M, Rahmstorf S (2011) Climate related sea-level variations over the past two millennia. *PNAS* 108(27):11017–11022
- King RS, Baker ME, Kazyak PF, Weller DE (2011) How novel is too novel? Stream community thresholds at exceptionally low levels of catchment urbanization. *Ecol Appl* 21(5):1659–1678. <https://doi.org/10.1890/10-1357.1>
- Kirwan ML, Gedan KB (2019) Sea-level driven land conversion and the formation of ghost forests. *Nat Clim Chang* 9(6):450–457. <https://doi.org/10.1038/s41558-019-0488-7>
- Kozłowski TT (1997) Responses of woody plants to flooding and salinity. *Tree physiology monograph* no. 1. Heron Publishing, Victoria, Canada, pp 1–29
- Krauss KW, Chambers JL, Allen JA (1998) Salinity effects and differential germination of several half-sib families of baldcypress from different seed sources. *New for* 15:53–68
- Manda AK, Giuliano AS, Allen TR (2014) Influence of artificial channels on the source and extent of saline water intrusion in the wind tide dominated wetlands of the southern Albemarle estuarine system (USA). *Environ Earth Sci* 71:4409–4419. <https://doi.org/10.1007/s12665-013-2834-9>
- Middleton BA (2016) Differences in impacts of Hurricane Sandy on freshwater swamps on the Delmarva Peninsula, Mid-Atlantic Coast, USA. *Ecol Eng* 87:62–70. <https://doi.org/10.1016/j.ecoleng.2015.11.035>
- Moleno GD, Defne Z, Aretxabaleta AL, Ganju NK, Carr JA (2021) Quantifying slopes as a driver of forest to marsh conversion using geospatial techniques: application to chesapeake bay coastal-plain, United States. *Front Environ Sci* 9:616319. <https://doi.org/10.3389/fenvs.2021.616319>
- Moorhead KK, Brinson MM (1995) Response of wetlands to rising sea level in the lower coastal plain of North Carolina. *Ecol Appl* 5(1):261–271. <https://doi.org/10.2307/1942068>
- Munns R, Tester M (2008) Mechanisms of salinity tolerance. *Annu Rev Plant Biol* 59(1):651–681. <https://doi.org/10.1146/annurev.arplant.59.032607.092911>
- Negrão S, Schmöckel SM, Tester M (2017) Evaluating physiological responses of plants to salinity stress. *Ann Bot* 119(1):1–11. <https://doi.org/10.1093/aob/mcw191>
- Noss RF, Platt WJ, Sorrie BA, Weakley AS, Means DB, Costanza J, Peet RK (2015) How global biodiversity hotspots may go unrecognized: lessons from the North American Coastal Plain. *Divers Distrib* 21(2):236–244. <https://doi.org/10.1111/ddi.12278>
- Osland MJ, Grace JB, Guntenspergen GR et al (2019) Climatic controls on the distribution of foundation plant species in coastal wetlands of the conterminous united states: knowledge gaps and emerging research needs. *Estuar Coasts* 42:1991–2003. <https://doi.org/10.1007/s12237-019-00640-z>

- Oksanen J, Blanchet FG, Kindt R, Legendre P, O'Hara RB, et al. (2011) vegan: community ecology package. R package version 1, pp 17–10
- Parida AK, Das AB (2005) Salt tolerance and salinity effects on plants: a review. *Ecotoxicol Environ Saf* 60(3):324–349. <https://doi.org/10.1016/j.ecoenv.2004.06.010>
- Parihar P, Singh S, Singh R, Singh VP, Prasad SM (2015) Effect of salinity stress on plants and its tolerance strategies: a review. *Environ Sci Pollut Res* 22(6):4056–4075. <https://doi.org/10.1007/s11356-014-3739-1>
- Peet RK, White PS, Wentworth TR (1998) A flexible, multi-purpose method for recording vegetation composition and structure. *Castanea* 63(3):262–274
- Pennings SC, Grant MB, Bertness MD (2005) Plant zonation in low-latitude salt marshes: disentangling the roles of flooding, salinity and competition. *J Ecol* 93(1):159–167. <https://doi.org/10.1111/j.1365-2745.2004.00959.x>
- Pezeshki SR, Delaune RD, Patrick WH (1990) Flooding and saltwater intrusion: potential effects on survival and productivity of wetland forests along the U.S. Gulf Coast. *For Ecol Manage* 33–34(C):287–301. [https://doi.org/10.1016/0378-1127\(90\)90199-L](https://doi.org/10.1016/0378-1127(90)90199-L)
- Poulter B (2005) Interactions between landscape disturbance and gradual environmental change: plant community migration in response to fire and sea level rise. Ph.D. Dissertation. Duke University. Durham, NC
- Poulter B, Christensen NL, Qian SS (2008) Tolerance of *Pinus taeda* and *Pinus serotina* to low salinity and flooding: implications for equilibrium vegetation dynamics. *J Veg Sci* 19(1):15–22. <https://doi.org/10.3170/2007-8-18410>
- Poulter B, Feldman RL, Brinson MM, Horton BP, Orbach MK, Pearsall SH, Reyes E, Riggs SR, Whitehead JC (2009a) Sea-level rise research and dialogue in North Carolina: creating windows for policy change. *Ocean Coast Manage* 52(3–4):147–153. <https://doi.org/10.1016/j.ocecoaman.2008.09.010>
- Poulter B, Qian SS, Christensen NL (2009b) Determinants of coastal treeline and the role of abiotic and biotic interactions. *Plant Ecol* 202(1):55–66. <https://doi.org/10.1007/s11258-008-9465-3>
- R Core Team (2019) R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>.
- Scheffer M, Carpenter SR, Lenton TM, Bascompte J, Brock W, Dakos V, van de Koppel J, van de Leemput IA, Levin SA, van Nes EH, Pascual M, Vandermeer J (2012) Anticipating critical transitions. *Science* 338(6105):334–338
- Schile LM, Callaway JC, Suding KN, Kelly NM (2017) Can community structure track sea-level rise? Stress and competitive controls in tidal wetlands. *Ecol Evol* 7(4):1276–1285. <https://doi.org/10.1002/ece3.2758>
- Schuerch M, Spencer T, Temmerman S, Kirwan ML, Wolff C, Lincke D, McOwen CJ, Pickering MD, Reef R, Vafeidis AT, Hinkel J, Nicholls RJ, Brown S (2018) Future response of global coastal wetlands to sea-level rise. *Nature* 561:231–234. <https://doi.org/10.1038/s41586-018-0476-5>
- Smart LS, Taillie PJ, Poulter B, Vukomanovic J, Singh KK, Swenson JJ, Helena Mitasova H, Smith JW, Meentemeyer RK (2020) Aboveground carbon loss associated with the spread of ghost forests as sea level rises. *Environ Res Lett*. <https://doi.org/10.1088/1748-9326/aba136>
- Sutter LA, Chambers RM, Perry JE (2015) Seawater intrusion mediates species transition in low salinity, tidal marsh vegetation. *Aquat Bot* 122:32–39. <https://doi.org/10.1016/j.aquabot.2015.01.002>
- Taillie PJ, Moorman CE, Poulter B, Ardón M, Emanuel RE (2019a) Decadal-scale vegetation change driven by salinity at leading edge of rising sea level. *Ecosystems* 22(8):1918–1930. <https://doi.org/10.1007/s10021-019-00382-w>
- Taillie PJ, Moorman CE, Smart LS, Pacifici K (2019b) Bird community shifts associated with saltwater exposure in coastal forests at the leading edge of rising sea level. *PLoS ONE* 14(5):e0216540
- Taillie P, Poulter B (2019) Eastern North Carolina coastal vegetation inventory. PANGAEA. <https://doi.org/10.1594/PANGAEA.896941>
- Tully KL, Weissman D, Wyner WJ, Miller J, Jordan T (2019) Soils in transition: saltwater intrusion alters soil chemistry in agricultural fields. *Biogeochemistry* 142(3):339–356. <https://doi.org/10.1007/s10533-019-00538-9>
- Ury EA, Anderson SM, Peet RK, Bernhardt ES, Wright JP (2020) Succession, regression and loss: does evidence of saltwater exposure explain recent changes in the tree communities of North Carolina's Coastal Plain? *Ann Bot*. <https://doi.org/10.1093/aob/mcz039/>
- Ury EA, Yang X, Wright JP, Bernhardt ES (2021) Rapid deforestation of a coastal landscape driven by sea-level rise and extreme events. *Ecol Appl*. <https://doi.org/10.1002/eap.2339>
- White EE, Ury EA, Bernhardt ES, Yang X (2021) Climate change driving widespread loss of coastal forested wetlands throughout the North American coastal plain. *Ecosystems*. <https://doi.org/10.1007/s10021-021-00686-w>
- Zhou M, Butterbach-Bahl K, Vereecken H, Brüggemann N (2017) A meta-analysis of soil salinization effects on nitrogen pools, cycles and fluxes in coastal ecosystems. *Glob Change Biol* 23(3):1338–1352. <https://doi.org/10.1111/gcb.13430>

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.